



**PCT**

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : <b>C12N 15/12, 15/85, C07K 15/06, 15/14, C12Q 1/68, G01N 33/53, A61K 39/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 94/23031</b> (43) International Publication Date: 13 October 1994 (13.10.94)
(21) International Application Number: <b>PCT/US94/02877</b> (22) International Filing Date: 17 March 1994 (17.03.94) (30) Priority Data: 08/037,230 26 March 1993 (26.03.93) <b>US</b> (71) Applicant: <b>LUDWIG INSTITUTE FOR CANCER RE- SEARCH [US/US]; 1345 Avenue of the Americas, New York, NY 10105 (US).</b> (72) Inventors: <b>GAUGLER, Beatrice; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE). VAN DEN EYNDE, Benoit; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE). BOON-FALLEUR, Thierry; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE). VAN DER BRUGGEN, Pierre; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE).</b> (74) Agent: <b>HANSON, Norman, D.; Felfe &amp; Lynch, 805 Third Avenue, New York, NY 10022 (US).</b>		(81) Designated States: <b>AU, CA, FI, JP, KR, NO, NZ, PT, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</b>  <b>Published</b> <i>With international search report.</i>	
(54) Title: <b>ISOLATED NUCLEIC ACID MOLECULES CODING FOR TUMOR REJECTION ANTIGEN PRECURSOR MAGE-3 AND USES THEREOF</b>			
(57) Abstract <p>The invention relates to nucleic acid molecules which code for the tumor rejection antigen precursor MAGE-3. Also disclosed are vectors, cell lines, and so forth, which utilize the nucleic acid molecule, and optionally, molecules coding for human leukocyte antigen HLA-A1. Uses of these materials in therapeutic and diagnostic contexts are also a part of the invention.</p>			

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

ISOLATED NUCLEIC ACID MOLECULES CODING FOR  
TUMOR REJECTION ANTIGEN PRECURSOR MAGE-3  
AND USES THEREOF

5     **RELATED APPLICATION**

          This application is a continuation-in-part of PCT Application PCT/US92/04354 filed on May 22, 1992 designating the United States, which is a continuation-in-part of Serial Number 807,043, filed December 12, 1991, which is a  
10     continuation-in-part of Serial Number 764,364, filed September 23, 1991, which is a continuation-in-part of Serial Number 728,838, filed July 9, 1991, which is a continuation-in-part of Serial Number 705,702, filed May 23, 1991, and now abandoned.

15

**FIELD OF THE INVENTION**

          This invention relates in general to the field of immunogenetics as applied to the study of oncology. More specifically, it relates to the study and analysis of  
20     mechanisms by which tumors are recognized by the organism's immune system such as through the presentation of so-called tumor rejection antigens, and the expression of what will be referred to herein as "tumor rejection antigen precursors" or "TRAPs". Most specifically, it refers to nucleic acid  
25     molecules coding for one such TRAP, i.e., MAGE-3, which is processed to a tumor rejection antigen or "TRA" presented by HLA-A1 molecules.

**BACKGROUND AND PRIOR ART**

30     The study of the recognition or lack of recognition of cancer cells by a host organism has proceeded in many different directions. Understanding of the field presumes some understanding of both basic immunology and oncology.

          Early research on mouse tumors revealed that these  
35     displayed molecules which led to rejection of tumor cells when transplanted into syngeneic animals. These molecules are "recognized" by T-cells in the recipient animal, and provoke

a cytolytic T-cell response with lysis of the transplanted cells. This evidence was first obtained with tumors induced in vitro by chemical carcinogens, such as methylcholanthrene. The antigens expressed by the tumors and which elicited the T-cell response were found to be different for each tumor. See  
5 Prehn, et al., J. Natl. Canc. Inst. 18: 769-778 (1957); Klein et al., Cancer Res. 20: 1561-1572 (1960); Gross, Cancer Res. 3: 326-333 (1943), Basombrio, Cancer Res. 30: 2458-2462 (1970) for general teachings on inducing tumors with chemical  
10 carcinogens and differences in cell surface antigens. This class of antigens has come to be known as "tumor specific transplantation antigens" or "TSTAs". Following the observation of the presentation of such antigens when induced by chemical carcinogens, similar results were obtained when  
15 tumors were induced in vitro via ultraviolet radiation. See Kripke, J. Natl. Canc. Inst. 53: 333-1336 (1974).

While T-cell mediated immune responses were observed for the types of tumor described supra, spontaneous tumors were thought to be generally non-immunogenic. These were therefore  
20 believed not to present antigens which provoked a response to the tumor in the tumor carrying subject. See Hewitt, et al., Brit. J. Cancer 33: 241-259 (1976).

The family of tum<sup>-</sup> antigen presenting cell lines are immunogenic variants obtained by mutagenesis of mouse tumor  
25 cells or cell lines, as described by Boon et al., J. Exp. Med. 152: 1184-1193 (1980), the disclosure of which is incorporated by reference. To elaborate, tum<sup>-</sup> antigens are obtained by mutating tumor cells which do not generate an immune response in syngeneic mice and will form tumors (i.e., "tum<sup>+</sup>" cells).  
30 When these tum<sup>-</sup> cells are mutagenized, they are rejected by syngeneic mice, and fail to form tumors (thus "tum<sup>-</sup>"). See Boon et al., Proc. Natl. Acad. Sci. USA 74: 272 (1977), the disclosure of which is incorporated by reference. Many tumor types have been shown to exhibit this phenomenon. See, e.g.,  
35 Frost et al., Cancer Res. 43: 125 (1983).

It appears that tum<sup>-</sup> variants fail to form progressive tumors because they elicit an immune rejection process. The

evidence in favor of this hypothesis includes the ability of "tum" variants of tumors, i.e., those which do not normally form tumors, to do so in mice with immune systems suppressed by sublethal irradiation, Van Pel et al., Proc. Natl. Acad. Sci. USA 76: 5282-5285 (1979); and the observation that intraperitoneally injected tum cells of mastocytoma P815 multiply exponentially for 12-15 days, and then are eliminated in only a few days in the midst of an influx of lymphocytes and macrophages (Uyttenhove et al., J. Exp. Med. 152: 1175-1183 (1980)). Further evidence includes the observation that mice acquire an immune memory which permits them to resist subsequent challenge to the same tum variant, even when immunosuppressive amounts of radiation are administered with the following challenge of cells (Boon et al., Proc. Natl. Acad. Sci. USA 74: 272-275 (1977); Van Pel et al., supra; Uyttenhove et al., supra). Later research found that when spontaneous tumors were subjected to mutagenesis, immunogenic variants were produced which did generate a response. Indeed, these variants were able to elicit an immune protective response against the original tumor. See Van Pel et al., J. Exp. Med. 157: 1992-2001 (1983). Thus, it has been shown that it is possible to elicit presentation of a so-called "tumor rejection antigen" in a tumor which is a target for a syngeneic rejection response. Similar results have been obtained when foreign genes have been transfected into spontaneous tumors. See Fearson et al., Cancer Res. 48: 2975-1980 (1988) in this regard.

A class of antigens has been recognized which are presented on the surface of tumor cells and are recognized by cytotoxic T cells, leading to lysis. This class of antigens will be referred to as "tumor rejection antigens" or "TRAs" hereafter. TRAs may or may not elicit antibody responses. The extent to which these antigens have been studied, has been via cytolytic T cell characterization studies, in vitro i.e., the study of the identification of the antigen by a particular cytolytic T cell ("CTL" hereafter) subset. The subset proliferates upon recognition of the presented tumor rejection

antigen, and the cells presenting the antigen are lysed. Characterization studies have identified CTL clones which specifically lyse cells expressing the antigens. Examples of this work may be found in Levy et al., Adv. Cancer Res. 24: 1-59 (1977); Boon et al., J. Exp. Med. 152: 1184-1193 (1980); Brunner et al., J. Immunol. 124: 1627-1634 (1980); Maryanski et al., Eur. J. Immunol. 124: 1627-1634 (1980); Maryanski et al., Eur. J. Immunol. 12: 406-412 (1982); Palladino et al., Canc. Res. 47: 5074-5079 (1987). This type of analysis is required for other types of antigens recognized by CTLs, including minor histocompatibility antigens, the male specific H-Y antigens, and a class of antigens, referred to as "tum-" antigens, and discussed herein.

A tumor exemplary of the subject matter described supra is known as P815. See DePlaen et al., Proc. Natl. Acad. Sci. USA 85: 2274-2278 (1988); Szikora et al., EMBO J 9: 1041-1050 (1990), and Sibille et al., J. Exp. Med. 172: 35-45 (1990), the disclosures of which are incorporated by reference. The P815 tumor is a mastocytoma, induced in a DBA/2 mouse with methylcholanthrene and cultured as both an in vitro tumor and a cell line. The P815 line has generated many tum<sup>-</sup> variants following mutagenesis, including variants referred to as P91A (DePlaen, supra), 35B (Szikora, supra), and P198 (Sibille, supra). In contrast to tumor rejection antigens - and this is a key distinction - the tum<sup>-</sup> antigens are only present after the tumor cells are mutagenized. Tumor rejection antigens are present on cells of a given tumor without mutagenesis. Hence, with reference to the literature, a cell line can be tum<sup>-</sup>, such as the line referred to as "P1", and can be provoked to produce tum<sup>-</sup> variants. Since the tum<sup>-</sup> phenotype differs from that of the parent cell line, one expects a difference in the DNA of tum<sup>-</sup> cell lines as compared to their tum<sup>-</sup> parental lines, and this difference can be exploited to locate the gene of interest in tum<sup>-</sup> cells. As a result, it was found that genes of tum<sup>-</sup> variants such as P91A, 35B and P198 differ from their normal alleles by point mutations in the coding regions of the gene. See Szikora and Sibille, supra, and Lurquin et

al., Cell 58: 293-303 (1989). This has proved not to be the case with the TRAs of this invention. These papers also demonstrated that peptides derived from the tum antigen are presented by the L<sup>d</sup> molecule for recognition by CTLs. P91A is presented by L<sup>d</sup>, P35 by D<sup>d</sup> and P198 by K<sup>d</sup>.

Prior patent applications PCT/US92/04354, U.S. Serial No. 807,043; 764,364; 728,838 and 707,702, all of which are incorporated by reference, describe inventions involving, inter alia, genes and other nucleic acid molecules which code for various TRAPs, which are in turn processed to tumor rejection antigen, or "TRAs".

The genes are useful as a source for the isolated and purified tumor rejection antigen precursor and the TRA themselves, either of which can be used as an agent for treating the cancer for which the antigen is a "marker", as well as in various diagnostic and surveillance approaches to oncology, discussed infra. It is known, for example, that tum cells can be used to generate CTLs which lyse cells presenting different tum antigens as well as tum cells. See, e.g., Maryanski et al., Eur. J. Immunol 12: 401 (1982); and Van den Eynde et al., Modern Trends in Leukemia IX (June 1990), the disclosures of which are incorporated by reference. The tumor rejection antigen precursor may be expressed in cells transfected by the gene, and then used to generate an immune response against a tumor of interest.

In the parallel case of human neoplasms, it has been observed that autologous mixed lymphocyte-tumor cell cultures ("MLTC" hereafter) frequently generate responder lymphocytes which lyse autologous tumor cells and do not lyse natural killer targets, autologous EBV-transformed B cells, or autologous fibroblasts (see Anichini et al., Immunol. Today 8: 385-389 (1987)). This response has been particularly well studied for melanomas, and MLTC have been carried out either with peripheral blood cells or with tumor infiltrating lymphocytes. Examples of the literature in this area including Knuth et al., Proc. Natl. Acad. Sci. USA 86: 2804-2802 (1984); Mukherji et al., J. Exp. Med. 158: 240 (1983);

Hérin et al., Int. J. Canc. 39: 390-396 (1987); Topalian et al., J. Clin. Oncol 6: 839-853 (1988). Stable cytotoxic T cell clones ("CTLs" hereafter) have been derived from MLTC responder cells, and these clones are specific for the tumor cells. See Mukherji et al., supra, Hérin et al., supra, Knuth et al., supra. The antigens recognized on tumor cells by these autologous CTLs do not appear to represent a cultural artifact, since they are found on fresh tumor cells. Topalian et al., supra; Degiovanni et al., Eur. J. Immunol. 20: 1865-1868 (1990). These observations, coupled with the techniques used herein to isolate the genes for specific murine tumor rejection antigen precursors, have led to the isolation of nucleic acid sequences coding for tumor rejection antigen precursors of TRAs presented on human tumors. It is now possible to isolate the nucleic acid sequences which code for tumor rejection antigen precursors, including, but not being limited to those most characteristic of a particular tumor, with ramifications that are described infra.

Additional work has focused upon the presentation of TRAs by the class of molecules known as human leukocyte antigens, or "HLAs". This work has resulted in several unexpected discoveries regarding the field. Specifically in U.S. patent application Serial Number 938,334, the disclosure of which is incorporated by reference, nonapeptides are taught which are presented by the HLA-A1 molecule. The reference teaches that given the known specificity of particular peptides for particular HLA molecules, one should expect a particular peptide to bind one HLA molecule, but not others. This is important, because different individuals possess different HLA phenotypes. As a result, while identification of a particular peptide as being a partner for a specific HLA molecule has diagnostic and therapeutic ramifications, these are only relevant for individuals with that particular HLA phenotype. There is a need for further work in the area, because cellular abnormalities are not restricted to one particular HLA phenotype, and targeted therapy requires some knowledge of the phenotype of the abnormal cells at issue.



In U.S. Patent Application Serial Number 008,446, filed January 22, 1993 and incorporated by reference, the fact that the MAGE-1 expression product is processed to a second TRA is disclosed. This second TRA is presented by HLA-C10-molecules. The disclosure shows that a given TRAP can yield a plurality of TRAs.

In U.S. Patent Application Serial Number 994,928, filed December 22, 1992, and incorporated by reference herein, tyrosinase is described as a tumor rejection antigen precursor. This reference discloses that a molecule which is produced by some normal cells (e.g., melanocytes), is processed in tumor cells to yield a tumor rejection antigen that is presented by HLA-A2 molecules.

It was mentioned, supra, that different individuals possess different HLA types. It has also been found that the expression of particular MAGE genes is not always linked to particular disorders, or individuals of particular HLA types. Thus, one cannot state, e.g., that all melanoma patients will express MAGE-1 TRAP nor could one say categorically that MAGE-1 expression is limited to melanoma patients of type HLA-A1. Further, one cannot state that only one type of TRAP is expressed in individuals of a particular HLA type. No rules or guidelines can be pointed to which correlate any of these factors.

Thus, it is not expected that a second TRAP is processed to a TRAP which is presented by HLA-A1 molecules. It has now been found that in addition to MAGE-1, a TRA derived from MAGE-3 TRAP is presented by HLA-A1 molecules. This is shown in examples 37-40, which follow, together with a discussion of the ramifications of this discovery.

These and various other aspects of the invention are elaborated upon in the disclosure which follows.

#### **BRIEF DESCRIPTION OF THE FIGURES**

Figure 1 depicts detection of transfectants expressing antigen P815A.

Figure 2 shows the sensitivity of clones P1.HTR, PO.HTR, genomic transfectant P1A.T2 and cosmid transfectant P1A.TC3.1

to lysis by various CTLs, as determined by chromium release assays.

Figure 3 is a restriction map of cosmid ClA.3.1.

5 Figure 4 shows Northern Blot analysis of expression of gene P1A.

Figure 5 sets forth the structure of gene P1A with its restriction sites.

10 Figure 6 shows the results obtained when cells were transfected with the gene from P1A, either isolated from P815 or normal cells and then tested with CTL lysis.

Figure 7 shows lytic studies using mast cell line L138.8A.

Figure 8 is a map of subfragments of the 2.4 kb antigen E fragment sequence which also express the antigen.

15 Figure 9 shows homology of sections of exon 3 from genes mage 1, 2 and 3.

Figure 10 shows the result of Northern blots for MAGE genes on various tissues.

20 Figure 11 presents the data of Figure 13 in table form. Figure 12 shows Southern Blot experiments using the various human melanoma cell lines employed in this application.

Figure 13 is a generalized schematic of the expression of MAGE 1, 2 and 3 genes by tumor and normal tissues.

25 Figure 14 shows results from a chromium release assay using CTL clone 20/38 on various cell lines.

Figure 15 presents the result of assays undertaken to determine antigenic specificity of CTL clone 20/38.

Figure 16 shows the results obtained when a TNF release assay was carried out on various transfected cells.

30 **BRIEF DESCRIPTION OF SEQUENCES**

SEQ ID NO: 1 is cDNA for part of gene P1A.

SEQ ID NO: 2 presents coding region of cDNA for gene P1A.

SEQ ID NO: 3 shows non coding DNA for P1A cDNA which is 3' to the coding region of SEQ ID NO: 2.

35 SEQ ID NO: 4 is the entire sequence of cDNA for P1A.

SEQ ID NO: 5 is the genomic DNA sequence for P1A.

SEQ ID NO: 6 shows the amino acid sequence for the

antigenic peptides for P1A TRA. The sequence is for cells which are A\* B\*, i.e., express both the A and B antigens.

SEQ ID NO: 7 is a nucleic acid sequence coding for antigen E.

5        SEQ ID NO: 8 is a nucleic acid sequence coding for MAGE-1.

SEQ ID NO: 9 is the gene for MAGE-2.

SEQ ID NO: 10 is the gene for MAGE-21.

SEQ ID NO: 11 is cDNA for MAGE-3.

10        SEQ ID NO: 12 is the gene for MAGE-31.

SEQ ID NO: 13 is the gene for MAGE-4.

SEQ ID NO: 14 is the gene for MAGE-41.

SEQ ID NO: 15 is cDNA for MAGE-4.

SEQ ID NO: 16 is cDNA for MAGE-5.

15        SEQ ID NO: 17 is genomic DNA for MAGE-51.

SEQ ID NO: 18 is cDNA for MAGE-6.

SEQ ID NO: 19 is genomic DNA for MAGE-7.

SEQ ID NO: 20 is genomic DNA for MAGE-8.

SEQ ID NO: 21 is genomic DNA for MAGE-9.

20        SEQ ID NO: 22 is genomic DNA for MAGE-10.

SEQ ID NO: 23 is genomic DNA for MAGE-11.

SEQ ID NO: 24 is genomic DNA for smage-I.

SEQ ID NO: 25 is genomic DNA for smage-II.

#### DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

25        Many different "MAGE" genes have been identified, as will be seen from the sequences which follow the application. The protocols described in the following examples were used to isolate these genes and cDNA sequences.

30        "MAGE" as used herein refers to a nucleic acid sequence isolated from human cells. The acronym "smage" is used to describe sequences of murine origin.

When "TRAP" or "TRAs" are discussed herein as being specific to a tumor type, this means that the molecule under consideration is associated with that type of tumor, although not necessarily to the exclusion of other tumor types.

35

#### Example 1

In order to identify and isolate the gene coding for

antigen P815A, gene transfection was used. This approach requires both a source of the gene, and a recipient cell line. Highly transfectable cell line P1.HTR was the starting material for the recipient, but it could not be used without further treatment, as it presents "antigen A", one of four recognized P815 tumor antigens. See Van Pel et al., Molecular Genetics 11: 467-475 (1985). Thus, screening experiments were carried out to isolate cell lines which did not express the antigen and which nonetheless possessed P1.HTR's desirable qualities.

To do this, P1.HTR was screened with CTLs which were specific for each of tumor antigens A, B, C and D. Such CTLs are described by Uyttenhove et al., J. Exp. Med. 157: 1040-1052 (1983).

To carry out the selection,  $10^6$  cells of P1.HTR were mixed with  $2-4 \times 10^6$  cells of the CTL clone in a round bottom tube in 2 ml of medium, and centrifuged for three minutes at 150xg. After four hours at 37°C, the cells were washed and resuspended in 10 ml of medium, following Maryanski et al., Eur. J. Immunol. 12: 406-412 (1982). Additional information on the CTL assay and screening protocol, in general may be found in Boon et al., J. Exp. Med. 152: 1184-1193 (1980), and Maryanski et al., Eur. J. Immunol. 12: 406-412 (1982), the disclosure of which are incorporated by reference.

When these selections were carried out, a cell line variant was found which expressed neither antigen A or B. Additional selections with CTLs specific for antigen C then yielded a variant which also lacked antigen C. Please see figure 2 for a summary of the results of these screenings. The variant PO.HTR is negative for antigens A, B and C, and was therefore chosen for the transfection experiments.

The cell line PO.HTR has been deposited in accordance with the Budapest Treaty at the Institute Pasteur Collection Nationale De Cultures De Microorganismes, 28, Rue de Docteur Roux, 75724 Paris France, and has accession number I-1117.

This methodology is adaptable to secure other cell lines which are variants of a cell type which normally presents at

least one of the four recognized P815 tumor antigens, i.e., antigens A, B, C and D, where the variants present none of antigens A, B and C. P1.HTR is a mastocytoma cell line, so it will be seen that the protocol enables the isolation of biologically pure mastocytoma cell lines which express none of P815 antigens A, B and C, but which are highly transfectable. Other tumor types may also be screened in this fashion to secure desired, biologically pure cell lines. The resulting cell lines should be at least as transfectable with foreign DNA as is P1.HTR, and should be selected so as to not express a specific antigen.

#### Example 2

Previous work reported by DePlaen et al., Proc. Natl. Acad. Sci. USA 85: 2274-2278 (1988) the disclosure of which is incorporated by reference herein had shown the efficacy of using cosmid library transfection to recover genes coding for tumor antigens.

Selective plasmid and genomic DNA of P1.HTR were prepared, following Wölfel et al., Immunogenetics 26: 178-187 (1987). The transfection procedure followed Corsaro et al., Somatic Cell Molec. Genet 7: 603-616 (1981), with some modification. Briefly, 60  $\mu$ g of cellular DNA and 3  $\mu$ g of DNA of plasmid pHMR272, described by Bernard et al., Exp. Cell. Biol. 158: 237-243 (1985) were mixed. This plasmid confers hygromycin resistance upon recipient cells, and therefore provides a convenient way to screen for transfectants. The mixed DNA was combined with 940  $\mu$ l of 1 mM Tris-HCl (pH 7.5), 0.1 mM EDTA; and 310  $\mu$ l 1M  $\text{CaCl}_2$ . The solution was added slowly, and under constant agitation to 1.25 ml of 50 mM Hepes, 280 mM NaCl, 1.5 mM  $\text{Na}_2\text{HPO}_4$ , adjusted to pH 7.1 with NaOH. Calcium phosphate - DNA precipitates were allowed to form for 30-45 minutes at room temperature. Following this, fifteen groups of P0.HTR cells ( $5 \times 10^6$ ) per group were centrifuged for 10 minutes at 400 g. Supernatants were removed, and pellets were resuspended directly into the medium containing the DNA precipitates. This mixture was incubated for 20 minutes at 37°C, after which it was added to

an 80 cm<sup>2</sup> tissue culture flask containing 22.5 ml DMEM, supplemented with 10% fetal calf serum. After 24 hours, medium was replaced. Forty-eight hours after transfection, cells were collected and counted. Transfected cells were selected in mass culture using culture medium supplemented with hygromycin B (350 ug/ml). This treatment selected cells for hygromycin resistance.

For each group, two flasks were prepared, each containing 8x10<sup>6</sup> cells in 40 ml of medium. In order to estimate the number of transfectants, 1x10<sup>6</sup> cells from each group were plated in 5 ml DMEM with 10% fetal calf serum (FCS), 0.4% bactoagar, and 300 ug/ml hygromycin B. The colonies were then counted 12 days later. Two independent determinations were carried out and the average taken. This was multiplied by 5 to estimate the number of transfectants in the corresponding group. Correction had to be made for the cloning efficiency of P815 cells, known to be about 0.3.

### Example 3

Eight days after transfection as described in example 2, supra, antibiotic resistant transfectants were separated from dead cells, using density centrifugation with Ficoll-Paque. These cells were maintained in non-selective medium for 1 or 2 days. The cells were plated in 96 well microplates (round bottom), at 30 cells/microwell in 200 ul of culture medium. Anywhere from 100-400 microwells were prepared, depending on the number of transfectants prepared. Agar colony tests gave estimates of 500-3000. After 5 days, the wells contained about 6x10<sup>4</sup> cells and replicate plates were prepared by transferring 1/10 of the wells to microplates which were then incubated at 30°C. One day later, master plates were centrifuged, medium removed, and 750 CTLs against P815 antigen A (CTL-P1:5) were added to each well together with 10<sup>6</sup> irradiated syngeneic feeder spleen cells in CTL culture medium containing 40 U/ml recombinant human IL-2, and HAT medium to kill stimulator cells. Six days later, plates were examined visually to identify wells where CTLs had proliferated. Where plates showed proliferating microcultures, aliquots of 100 ul

of the wells were transferred to another plate containing  $^{51}\text{Cr}$  labeled P1.HTR target cells ( $2 \times 10^3$  -  $4 \times 10^3$  per well), and chromium release was measured after 4 hours. Replicate microcultures corresponding to those showing high CTL activity were expanded and cloned by limited dilution in DMEM with 10% FCS. Five days later, about 200 clones were collected and screened with the CTL.P1:5 cell line, described supra, in a visual lysis assay. See figure 1A for these results.

In these experiments, three of the fifteen groups of transfectants yielded a few positive microcultures. These microcultures were tested for lytic activity against P1.HTR, as described supra. Most of the microcultures where proliferation was observed showed lytic activity. This activity was well above background, as shown in figure 1B. This figure summarizes data wherein two groups of cells (groups "5" and "14"), 400 and 300 microwells were seeded with 30 hygromycin resistant transfected cells. Amplification and duplication of the microcultures was followed by addition of anti-A CTL P1:5. Six days later, lytic activity against P1.HTR was tested. In the figure, each point represents lytic activity of a single microculture.

Duplicate microcultures corresponding to several positive wells were subcloned, and more than 1% of the subclones were found to be lysed by anti-A CTL. Thus, three independent transfectants expressing P815A were obtained from 33,000 hygromycin resistant transfectants. One of these lines, referred to hereafter as P1A.T2 was tested further.

The relevant antigen profile of P1A.T2 is shown in figure 2, this being obtained via anti-CTL assays of the type described supra.

#### Example 4

The CTL assays carried out for P1A.T2 demonstrated that it presented antigen A ("P815A"), and therefore had received the gene from P1.HTR. To that end, this cell line was used as a source for the gene for the antigen precursor in the following experiments.

Prior work had shown that genes coding for tum antigens

could be recovered directly from transfectants obtained with a cosmid library. See DePlaen et al., Proc. Natl. Acad. Sci. USA 85: 2274-2278 (1988). This procedure was followed for recovery of the P815 gene.

5       Total genomic DNA of P1A.T2 was partially digested with restriction endonuclease Sau 3A1, and fractionated by NaCl density gradient ultracentrifugation to enrich for 35-50 kb DNA fragments, following Grosveld et al., Gene 10: 6715-6732 (1982). These fragments were ligated to cosmid arms of C2RB, 10 described by Bates et al., Gene 26: 137-146 (1983), the disclosure of which is incorporated by reference. These cosmid arms had been obtained by cleavage with SmaI and treatment with calf intestinal phosphatase, followed by digestion with BamHI. Ligated DNA was packaged into lambda 15 phage components, and titrated on E. coli ED 8767, following Grosveld et al., supra. Approximately  $9 \times 10^5$  ampicillin resistant colonies were obtained per microgram of DNA insert.

The cosmid groups were amplified by mixing 30,000 independent cosmids with 2 ml of ED 8767 in 10 mM  $MgCl_2$ , 20 incubated 20 minutes at 37°C, diluted with 20 ml of Luria Bertani ("LB") medium, followed by incubation for one hour. This suspension was titrated and used to inoculate 1 liter of LB medium in the presence of ampicillin (50 ug/ml). At a 25 bacterial concentration of  $2 \times 10^8$  cells/ml ( $OD_{600}=0.8$ ), a 10 ml aliquot was frozen, and 200 ug/ml chloramphenicol was added to the culture for overnight incubation. Total cosmid DNA was isolated by alkaline lysis procedure, and purified on CsCl gradient.

In these experiments, a library of 650,000 cosmids was 30 prepared. The amplification protocol involved the use of 21 groups of approximately 30,000 cosmids.

#### Example 5

Using the twenty-one groups of cosmids alluded to supra, (60 ug) and 4 ug of pHMR272, described supra, groups of  $5 \times 10^6$  35 PO.HTR cells were used as transfectant hosts. Transfection was carried out in the same manner as described in the preceding experiments. An average of 3000 transfectants per



group were tested for antigen presentation, again using CTL assays as described. One group of cosmids repeatedly yielded positive transfectants, at a frequency of about 1/5,000 drug resistant transfectants. The transfectants, as with P1A.T2, also showed expression of both antigen A and B. The pattern of expression of transfectant P1A.TC3.1 is shown in figure 2.

#### Example 6

As indicated in Example 5, supra, three independent cosmid transfected cells presenting P815A antigen were isolated. The DNA of these transfectants was isolated and packaged directly with lambda phage extracts, following DePlaen et al., Proc. Natl. Acad. Sci. USA 85: 2274-2278 (1988). The resulting product was titrated on E. coli ED 8767 with ampicillin selection, as in Example 5. Similarly, amplification of the cosmids and transfection followed Example 5, again using PO.HTR.

High frequencies of transfection were observed, as described in Table 1, which follows:

**Table 1.** Transfer of the expression of antigen P815A by cosmids obtained by direct packaging

Transfectant obtained with the cosmid library	No. of cosmids obtained by direct packaging of 0.5 µg of DNA	No. of transfectants expressing P815A / no. of HmB <sup>r</sup> transfectants
TC3.1	32	87/192
TC3.2	32000	49/384
TC3.3	44	25/72

The cosmids were analyzed with restriction enzymes and it was found that directly packaged transfectant P1A.TC3.1 contained 32 cosmids, 7 of which were different. Each of these 7 cosmids was transfected into PO.HTR, in the manner described supra, and again, following the protocols described above, transfectants were studied for presentation of P815A.

Four of the cosmid transfectants showed P815A presentation and, as with all experiments described herein, P815B was co-expressed.

5 Of the four cosmids showing presentation of the two antigens, cosmid C1A.3.1 was only 16.7 kilobases long, and was selected for further analysis as described infra.

The cosmid C1A.3.1 was subjected to restriction endonuclease analysis, yielding the map shown in Figure 3.

10 All EcoRI fragments were transfected, again using the above described protocols, and only the 7.4 kilobase fragment produced a transfectant that anti-A CTLs could lyse. Similar experiments were carried out on the PstI fragments, and only a 4.1 kb fragment fully contained within the 7.4 kb EcoRI fragment produced lysable transfectants.

15 This fragment (i.e., the 4.1 kb PstI fragment), was digested with SmaI, giving a 2.3 kb fragment which also yielded host cells presenting antigens A and B after transfection. Finally, a fragment 900 bases long, secured with SmaI/XbaI, also transferred expression of the precursors  
20 of these two antigens, i.e., the transfected host cell presented both antigen A and antigen B.

#### Example 7

The 900 base fragment described above was used as a probe to detect the expression of the P815A gene in parent cell line  
25 P1.HTR. To accomplish this, total cellular RNA was first isolated using the guanidine-isothiocyanate procedure of Davis et al., Basic Methods In Molecular Biology (Elsevier Science Publishing Co, New York) (1986). The same reference was the source of the method used to  
30 isolate and purify polyA<sup>+</sup> mRNA using oligodT cellulose column chromatography.

Samples were then subjected to Northern Blot analysis. RNA samples were fractionated on 1% agarose gels containing 0.66 M formaldehyde. The gels were treated with 10xSSC (SSC:  
35 0.15 M NaCl; 0.015 M sodium citrate, pH 7.0) for 30 minutes before overnight blotting on nitrocellulose membranes. These were baked for two hours at 80°C, after which the membranes

were prehybridized for 15 minutes at 60°C in a solution containing 10% dextran sulfate, 1% SDS and 1M NaCl. Hybridization was then carried out using denatured probe (the 900 base fragment), together with 100 ug/ml salmon sperm DNA.

When this protocol was carried out using P1.HTR poly A<sup>+</sup> RNA, a band of 1.2 kb and two fainter bands were identified, as shown in Figure 4, lane 1 (6 ug of the RNA).

The same probe was used to screen a cDNA library, prepared from poly-A<sup>+</sup> RNA from the cell line. This yielded a clone with a 1kb insert, suggesting a missing 5' end. The Northern blots for the cDNA are not shown.

Hybridization experiments in each case were carried out overnight at 60°C. The blots were washed twice at room temperature with 2xSSC and twice at 60°C with 2xSSC supplemented with 1% SDS.

The foregoing experiments delineated the DNA expressing the P815A antigen precursor sufficiently to allow sequencing, using the well known Sanger dideoxy chain termination method. This was carried out on clones generated using a variety of restriction endonucleases and by specific priming with synthetic oligonucleotide primers. The results for exons of the gene are set forth in sequence id no: 4.

#### Example 8

The Northern analysis described supra suggested that the 5' end of the cDNA was missing. To obtain this sequence, cDNA was prepared from P1.HTR RNA using a primer corresponding to positions 320-303. The sequence was then amplified using the polymerase chain reaction using a 3' primer corresponding to positions 286-266 and a 5' primer described by Frohman et al., Proc. Natl. Acad. Sci. USA 85: 8998-9002 (1988). A band of the expected size (270 bases) was found, which hybridized to the 900 bp SmaI/XbaI fragment described supra on a Southern blot. Following cloning into ml3tg 130 λ tg 131, the small, 270 bp fragment was sequenced. The sequence is shown in sequence id no: 1.

#### Example 9

Following the procurement of the sequences described in Examples 7 and 8 and depicted in seq id no: 4, a 5.7 kb region of cosmid C1A.3.1 was sequenced. This fragment was known to contain the 900 base fragment which expressed P815A in transfectants. This experiment permitted delineation of introns and exons, since the cosmid is genomic in origin.

The delineated structure of the gene is shown in figure 5. Together with seq id no: 4, these data show that the gene for the antigen precursor, referred to as "PlA" hereafter, is approximately 5 kilobases long and contains 3 exons. An ORF for a protein of 224 amino acids starts in exon 1, ending in exon 2. The 900 base pair fragment which transfers expression of precursors for antigens A and B only contains exon 1. The promoter region contains a CAAT box, as indicated in seq. id no: 1, and an enhancer sequence. This latter feature has been observed in promoters of most MHC class I genes, as observed by Geraghty et al., J. Exp. Med 171: 1-18 (1990); Kimura et al., Cell 44: 261-272 (1986).

A computer homology search was carried out, using program FASTA with K-triple parameters of 3 and 6, as suggested by Lipman et al., Science 227: 1435-1441 (1985), and using Genbank database release 65 (October 1990). No homology was found except for a stretch of 95 bases corresponding to part of an acid region coded by exon 1 (positions 524-618), which is similar to sequences coding for acidic regions in mouse nucleolar protein NO38/B23, as described by Bourbon et al., Mol. Biol. 200: 627-638 (1988), and Schmidt-Zachmann et al., Chromosoma 96: 417-426 (1988). Fifty six of 95 bases were identical. In order to test whether these homologies were the reason for cross hybridizing, experiments were carried out using a mouse spleen cDNA library screened with the 900 base fragment. cDNA clones corresponding closely to the sizes of the cross hybridizing bands were obtained. These were partially sequenced, and the 2.6 kb cDNA was found to correspond exactly to reported cDNA sequence of mouse nucleolin, while the 1.5 kb cDNA corresponded to mouse nucleolar protein NO38/B23.

Analysis of the nucleotide sequence of the gene, referred to as "P1A" hereafter, suggests that its coded product has a molecular mass of 25 kd. Analysis of the sequence id no: 4 shows a potential nuclear targeting signal at residues 5-9 (Dingwall et al., Ann. Rev. Cell Biol. 2: 367-390 (1986)), as well as a large acidic domain at positions 83-118. As indicated supra, this contains the region of homology between P1A and the two nucleolar proteins. A putative phosphorylation site can be found at position 125 (serine). Also, a second acidic domain is found close to the C-terminus as an uninterrupted stretch of 14 glutamate residues. A similar C-terminal structure has been found by Kessel et al. Proc. Natl. Acad. Sci. USA 84: 5306-5310 (1987), in a murine homeodomain protein having nuclear localization.

In studies comparing the sequence of gene P1A to the sequences for P91A, 35B and P198, no similarities were found, showing that P1A is indicative of a different class of genes and antigens.

#### Example 10

With the P1A probe and sequence in hand, investigations were carried out to determine whether the gene present in normal tissue was identical to that expressed by the tumor. To do this, phage libraries were prepared, using lambda zapII 10 and genomic DNA of DBA2 murine kidney cells. P1A was used as a probe. Hybridization conditions were as described supra, and a hybridizing clone was found. The clone contained exons one and two of the P1A gene, and corresponded to positions - 0.7 to 3.8 of figure 5. Following localization of this sequence, PCR amplification was carried out to obtain the sequence corresponding to 3.8 to 4.5 of figure 5.

Sequence analysis was carried out, and no differences were found between the gene from normal kidneys and the P1A gene as obtained from the P815 tumor cells.

In further experiments, the gene as found in DBA/2 kidney cells was transfected into PO.HTR, as described supra. These experiments, presented pictorially in figure 7, showed that antigens A and B were expressed as efficiently by the kidney

gene isolated from normal kidney cells as with the P1A gene isolated from normal kidney cells.

These experiments lead to the conclusion that the gene coding for the tumor rejection antigen precursor is a gene that does not result from a mutation; rather, it would appear that the gene is the same as one present in normal cells, but is not expressed therein. The ramifications of this finding are important, and are discussed infra.

In studies not elaborated upon herein, it was found that variants of the gene were available. Some cells were "P1A-B+", rather than the normal "P1A". The only difference between these is a point mutation in exon 1, with the 18th triplet coding for Ala in the variant instead of Val.

#### Example 11

Additional experiments were carried out with other cell types. Following the protocols described for Northern blot hybridizations supra, RNA of normal liver and spleen cells was tested to determine if a transcript of the P1A gene could be found. The Northern blot data are presented in figure 4 and, as can be seen, there is no evidence of expression.

The murine P815 cell line from which P1A was isolated is a mastocytoma. Therefore, mast cell lines were studied to determine if they expressed the gene. Mast cell line MC/9, described by Nabel et al., Cell 23: 19-28 (1981), and short term cultures of bone marrow derived mast cells were tested in the manner described supra (Northern blotting), but no transcript was found. In contrast when a Balb/C derived IL-3 dependent cell line L138.8A (Hültner et al., J. Immunol. 142: 3440-3446 (1989)) was tested, a strong signal was found. The mast cell work is shown in figure 4.

It is known that both BALB/C and DBA/2 mice share H-2<sup>d</sup> haplotype, and thus it was possible to test sensitivity to lysis using the CTLs described supra. Figure 8 shows these results, which essentially prove that anti-A and anti-B CTLs lysed the cells strongly, whereas anti-C and anti-D lines did not.

Further tests were carried out on other murine tumor cell

lines, i.e., teratocarcinoma cell line PCC4 (Boon et al., Proc. Natl. Acad. Sci. USA 74: 272-275 (1977), and leukemias LEC and WEH1-3B. Expression could not be detected in any of these samples.

#### Example 12

The actual presentation of the P1A antigen by MHC molecules was of interest. To test this, cosmid C1A.3.1 was transfected into fibroblast cell line DAP, which shows phenotype H-2<sup>k</sup>. The cell lines were transfected with genes expressing one of the K<sup>d</sup>, D<sup>d</sup>, and L<sup>d</sup> antigen. Following transfection with both the cosmid and the MHC gene, lysis with CTLs was studied, again as described supra. These studies, summarized in Table 2, show that L<sup>d</sup> is required for presentation of the P1A antigens A and B.

Table 2. H-2-restriction of antigens P815A and P815B

Recipient cell*	No. of clones lysed by the CTL/ no. of HmB <sup>r</sup> clones*	
	CTL anti-A	CTL anti-B
DAP (H-2 <sup>k</sup> )	0/208	0/194
DAP + K <sup>d</sup>	0/165	0/162
DAP + D <sup>d</sup>	0/157	0/129
DAP + L <sup>d</sup>	25/33	15/20

\*Cosmid C1A.3.1 containing the entire P1A gene was transfected in DAP cells previously transfected with H-2<sup>d</sup> class I genes as indicated.

\*Independent drug-resistant colonies were tested for lysis by anti-A or anti-B CTL in a visual assay.

The observation that one may associate presentation of a tumor rejection antigen with a particular MHC molecule was confirmed in experiments with human cells and HLA molecules, as elaborated upon infra.

5     Example 13

Using the sequence of the P1A gene as well as the amino acid sequence derivable therefrom, antigenic peptides which were A\* B\* (i.e., characteristic of cells which express both the A and B antigens), and those which are A\*B\* were identified. The peptide is presented in Figure 10. This peptide when administered to samples of PO.HTR cells in the presence of CTL cell lines specific to cells presenting it, led to lysis of the PO.HTR cells, lending support to the view that peptides based on the product expressed by the gene can be used as vaccines.

15     Example 14

The human melanoma cell line referred to hereafter as MZ2-MEL is not a clonal cell line. It expresses four stable antigens recognized by autologous CTLs, known as antigens "D, E, F, and A". In addition, two other antigens "B" and "C" are expressed by some sublines of the tumor. CTL clones specific for these six antigens are described by Van den Eynde et al., Int. J. Canc. 44: 634-640 (1989). Among the recognized subclones of MZ2-MEL are MEL.43, MEL3.0 and MEL3.1. (Van den Eynde et al., supra). Cell line MEL3.1 expresses antigen E, as determined by CTL studies as described for P815 variants, supra, so it was chosen as a source for the nucleic acid sequence expressing the antigen precursor.

In isolating the pertinent nucleic acid sequence for a tumor rejection antigen precursor, the techniques developed supra, showed that a recipient cell is needed which fulfills two criteria: (i) the recipient cell must not express the TRAP of interest under normal conditions, and (ii) it must express the relevant class I HLA molecule. Also, the recipient cell must have a high transfection frequency, i.e., it must be a "good" recipient.

In order to secure such a cell line, the clonal subline



ME3.1 was subjected to repeated selection with anti-E CTL 82/30 as described by Van den Eynde, supra. The repeated cycles of selection led to isolation of subclone MZ2-MEL-2.2 isc E'. This subclone is also HPRT<sup>r</sup>, (i.e., sensitive to HAT medium:  $10^{-4}$  M hypoxanthine,  $3.8 \times 10^{-7}$  aminopterin,  $1.6 \times 10^{-5}$  M 2-deoxythymidine). The subclone is referred to as "MEL-2.2" for simplicity hereafter.

#### Example 15

The genomic DNA of MEL3.0 was prepared following Wölfel et al., Immunogenetics 26: 178-187 (1987), the disclosure of which is incorporated by reference. The plasmid pSVtkneo $\beta$ , as described by Nicolas et al., Cold Spring Harb., Conf. Cell Prolif. 10: 469-485 (1983) confers geneticin resistance, so it can be used as a marker for cotransfection, as it was in this experiment.

Following a procedure similar but not identical to that of Corsao et al., Somatic Cell Molec. Genet 7: 603-616 (1981), total genomic DNA and the plasmid were cotransfected. The genomic DNA (60  $\mu$ g) and plasmid DNA (6  $\mu$ g) were mixed in 940  $\mu$ l of 1 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, after which 310  $\mu$ l of 1M CaCl<sub>2</sub> was added. This solution was slowly added, under constant agitation, to 1.25 ml of 2xHBS (50 mM HEPES, 280 mM NaCl 1.5 mM Na<sub>2</sub>HPO<sub>4</sub>, adjusted to pH 7.1 with NaOH). The calcium phosphate DNA precipitates were allowed to form for 30-45 minutes at room temperature, after which they were applied to 80 cm<sup>2</sup> tissue culture flasks which had been seeded 24 hours previously with  $3 \times 10^6$  MEL2.2 cells, in 22.5 ml of melanoma culture medium (Dulbecco's Modified Eagle's Medium) supplemented with 10% fetal calf serum. After 24 hours, the medium was replaced. Forty eight hours after transfection, the cells were harvested and seeded at  $4 \times 10^6$  cells per 80 cm<sup>2</sup> flask in melanoma culture medium supplemented with 2 mg/ml of geneticin. The geneticin serves as a selection marker.

#### Example 16

Thirteen days after transfection, geneticin-resistant colonies were counted, harvested, and cultured in nonselective

medium for 2 or 3 days. Transfected cells were then plated in 96-well microplates at 200 cells/well in 200  $\mu$ l of culture medium with 20% fetal calf serum (FCS) in order to obtain approximately 30 growing colonies per well. The number of microcultures was aimed at achieving redundancy, i.e., such that every independent transfectant should be represented at least four times.

After 10 days, wells contained approximately  $6 \times 10^4$  cells. These cells were detached, and 1/3 of each microculture was transferred to a duplicate plate. After 6 hours, i.e., after readherence, medium was removed and 1500 anti-E CTL (CTL 82/30), were added to each well in 100  $\mu$ l of CTL culture medium with 35 U/ml of IL-2. One day later, the supernatant (50  $\mu$ l) was harvested and examined for TNF concentration, for reasons set forth in the following example.

#### Example 17

The size of the mammalian genome is  $6 \times 10^6$  kb. As the average amount of DNA integrated in each drug-resistant transfectant was expected to be about 200 kb, a minimum of 30,000 transfectants would need to be examined to ascertain whether antigen E had been transfected. Prior work with murine cells had shown that when a CTL stimulation assay was used, groups containing only 3% of cells expressing the antigen of interest could be identified. This should reduce the number of assays by a factor of 30. While an anti-E CTL assay, as described supra, in mixed E<sup>+</sup>/E<sup>-</sup> cells was helpful, it was not sufficient in that consistent results could not be obtained.

As a result, an alternative test was devised. Stimulation of CTLs was studied by release of tumor necrosis factor ("TNF") using well known methodologies which need not be repeated here. As described in Example 15, 1500 CTL 82/30 cells had been added per well of transfectants. These CTLs were collected 6 days after stimulation. As indicated supra, after 1/3 of the cells in each well had been removed and the remaining 2/3 ( $4 \times 10^4$ ) had readhered, the CTLs and IL-2 were added thereto. The 50  $\mu$ l of supernatant was removed 24 hours

later and transferred to a microplate containing  $3 \times 10^4$  W13 (WEHI-164 clone 13; Espevik et al., J. Immunol. Meth. 95: 99-105 (1986)) cells in 50  $\mu$ l of W13 culture medium (RPMI-1640, supplemented with L-arginine (116 mg/l), L-asparagine (36 mg/l), L-glutamine (216 mg/l), and 10% FCS supplemented with 2  $\mu$ g of actinomycin D at 37% in an 8% CO<sub>2</sub> atmosphere. The cell line W13 is a mouse fibrosarcoma cell line sensitive to TNF. Dilutions of recombinant TNF- $\beta$  in RPMI 1640 were added to target cell controls.

The W13 cultures were evaluated after 20 hours of incubation, and dead cell percentage was measured using an adaptation of the colorimetric assay of Hansen et al., J. Immunol. Meth. 119: 203-210 (1989). This involved adding 50  $\mu$ l of (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide at 2.5 mg/ml in PBS, followed by two hours of incubation at 37°C. Dark blue formazan crystals were dissolved by adding 100  $\mu$ l of lysis solution (1 volume N,N dimethyl formamide mixed at 37°C with two volumes of water containing 30% (w/v) sodium dodecyl sulphate, at pH 4.7 from 1.6% acetic acid and 2.5% 1N HCl). Plates were incubated at 37°C overnight, and ODs were taken at 570 nm using 650 nm as control. Dead cell percentage was determined via the formula:

$$100 \times \left[ 1 - \frac{100 - (\text{OD}_{570} \text{ sample well})}{\text{OD}_{570} \text{ well} + \text{medium}} \right]$$

following Espevik et al., J. Immunol. Meth. 95: 99-105 (1986). The results showed that even when the ratio of E<sup>+</sup>/E<sup>-</sup> cells was as low as 1/45, significant production of TNF was observed, thus showing active CTLs. This led to the decision to test the drug resistant transfectants in groups of 30.

#### Example 18

Cells were tested for TNF production as discussed in Example 17, supra. A total of 100 groups of E<sup>-</sup> cells ( $4 \times 10^6$  cells/group) were tested following transfection, and  $7 \times 10^4$  independent geneticin resistant transfectants were obtained, for an average of 700 per group. Only one group of

transfected cells led to a microculture which caused anti-E antigen CTL clone 82/30 to produce TNF. Of 300 clones tested, 8 were positive. These clones were then tested for lysis by anti-E CTL, using the standard  $^{51}\text{Cr}$  release assay, and were found to be lysed as efficiently as the original E<sup>+</sup> cell line. The transfectant E.T1, discussed herein, had the same lysis pattern as did MEL2.2 for CTLs against antigens B,C,D and F.

The fact that only one transfectant presented the antigen out of 70,000 geneticin resistance transfectants may at first seem very low, but it is not. The work described supra for P815 showed an average frequency of 1/13,000. Human DNA recipient MEL2.2 appears to integrate 5 times less DNA than P1.HTR.

#### Example 19

Once transfectant E.T1 was found, analysis had to address several questions including whether an E<sup>+</sup> contaminant of the cell population was the cause. The analysis of antigen presentation, described supra, shows that E.T1 is B<sup>+</sup> and C<sup>+</sup>, just like the recipient cell MEL2.2. It was also found to be HPRT<sup>+</sup>, using standard selection procedures. All E<sup>+</sup> cells used in the work described herein, however, were HPRT<sup>+</sup>.

It was also possible that an E<sup>+</sup> revertant of MEL2.2 was the source for E.T1. To test this, the observation by Perucho et al., Cell 22: 309-317 (1980), that cotransfected sequences usually integrate together at a single location of recipient genome was employed. If antigen E in a transfectant results from cotransfection with pSVtkneo $\beta$ , then sequences should be linked and deletion of the antigen might also delete the neighboring pSVtkneo $\beta$  sequences. Wölfel et al., supra, has shown this to be true. If a normally E<sup>+</sup> cell is transfected with pSVtkneo $\beta$ , then sequences should be linked and deletion of the antigen might also delete the neighboring pSVtkneo $\beta$  sequences. If a normally E<sup>+</sup> cell transfected with pSVtkneo $\beta$  is E.T1, however, "co-deletion" should not take place. To test this, the transfectant E.T1 was subjected to immunoselection with 82/30, as described supra. Two antigen loss variants were obtained, which resisted lysis by this CTL.

Neither of these had lost geneticin resistance; however, Southern blot analysis showed loss of several *neo<sup>r</sup>* sequences in the variants, showing close linkage between the E gene and *neo<sup>r</sup>* gene in E.T1, leading to the conclusion that E.T1 was a transfectant.

#### Example 20

The E<sup>+</sup> subclone MZ2-MEL 4B was used as a source of DNA for preparation of a cosmid library. This library of nearly 700,000 cosmids was transfected into MZ2-MEL 2.2 cells, following the cosmid transfection protocols described supra.

By packaging the DNA of cosmid transfectants directly into lambda phase components, it is sometimes possible to retrieve cosmids that contain the sequences of interest. This procedure was unsuccessful here, so we rescued the transfected sequence by ligating DNA of the transfectant to appropriate restriction fragments of cosmid vector pTL6. This was tried with two transfectants and was successful with one of them. One cosmid, referred to as B3, was recovered from this experiment, and subjected to restriction endonuclease digestion via XmaI, or by BamHI digestion of a large, 12 kb XmaI transfected fragment. The fragments were cloned into vector pTZ 18R, and then transfected into MEL2.2. Again, TNF production was the measure by which successful transfection was determined. The experiments led to the determination of a gene sequence capable of transfecting antigen E on the 12 kb XmaI fragment, and then on the 2.4 kb fragment of BamHI digestion of the 12 kb segment.

The 2.4 kb fragment hybridizes with a 2.4 kb fragment from MZ2-MEL and with a T cell clone of patient MZ-2, as determined by Southern Blots (BamHI/SmaI digested DNA). The band is absent from E<sup>-</sup> antigen loss variants of MZ2-MEL, as seen in Figure 12.

The sequence for the E antigen precursor gene has been determined, and is presented herein:

	10	20	30	40	50	60
1	GGATCCAGGC	CCTGCCAGGA	AAAATATAAG	GGCCCTGCCGT	GAGAACAGAG	GGGGTCATCC 60
61	ACTGCCATGAG	AGTGGGGATG	TCACAGAGTC	CAGCCCACCC	TCCTGGTAGC	ACTGAGAAGC 120
121	CAGGGCTGTG	CTTGCGGTCT	GCACCCTGAG	GGCCCGTGGA	TTCTCTTTCC	TGGAGCTCCA 180
181	GGAACCAGGC	AGTGAGGCCT	TGGTCTGAGA	CAGTATCCTC	AGGTCACAGA	GCAGAGGATG 240
241	CACAGGGTGT	GCCAGCAGTG	AATGTTTGCC	CTGAATGCAC	ACCAAGGGCC	CCACCTGCCA 300
301	CAGGACACAT	AGGACTCCAC	AGAGTCTGGC	CTCACCTCCC	TACTGTCACT	CCTGTAGAAT 360
361	CGACCTCTGC	TGGCCGGCTG	TACCCTGAGT	ACCCTCTCAC	TTCTCTCTTC	AGGTTTTTCAG 420
421	GGGACAGGCC	AACCCAGAGG	ACAGGATTCC	CTGGAGGGCA	CAGAGGAGCA	CCAAGGAGAA 480
481	GATCTGTAAG	TAGGCCCTTG	TTAGAGTCTC	CAAGGTTTCT	TTCTCAGCTG	AGGCCTCTCA 540
541	CACACTCCCT	CTCTCCCCAG	GCCTGTGGGT	CTTCATTGCC	CAGCTCCTGC	CCACACTCCT 600
601	GCCTGTGCCC	CTGACGAGAG	TCATCATGTC	TCTTGAGCAG	AGGAGTCTGC	ACTGCAAGCC 660
661	TGAGGAAGCC	CTTGAGGCCC	AACAAGAGGC	CCTGGGCTCG	GTGTGTGTGC	AGGCTGCCAC 720
721	CTCCTCTCTC	TCTCCTCTGG	TCCTGGGCAC	CCTGGAGGAG	GTGCCCACTG	CTGGGTCAAC 780
781	AGATCCTCCC	CAGAGTCCTC	AGGGAGCCTC	CGCCTTTCCC	ACTACCATCA	ACTTCACTCG 840
841	ACAGAGGCAA	CCCAGTGAGG	GTTCCAGCAG	CCGTGAAGAG	GAGGGGCCAA	GCACCTCTTG 900
901	TATCCTGGAG	TCCTTGTTC	GAGCAGTAAT	CACTAAGAAG	GTGGCTGATT	TGGTTGGTTT 960
961	TCTGCTCCTC	AAATATCGAG	CCAGGGAGCC	AGTCACAAAG	GCAGAAATGC	TGGAGAGTGT 1020
1021	CATCAAAAAT	TACAAGCACT	GTTTTCTGTA	GATCTTCGGC	AAAGCCTCTG	AGTCCTTGCA 1080
1081	GCTGGTCTTT	GGCATTGACG	TGAAGGAAGC	AGACCCACC	GGCCACTCCT	ATGTCTCTGT 1140
1141	CACCTGCCTA	GGTCTCTCCT	ATGATGGCCT	GCTGGGTGAT	AATCAGATCA	TGCCCAAGAC 1200
1201	AGGCTTCCTG	ATAATTGTCC	TGGTCATGAT	TGCAATGGAG	GGCGGCCATG	CTCCTGAGGA 1260
1261	GGAATCTGG	GAGGAGCTGA	GTGTGATGGA	GGTGTATGAT	GGGAGGGAGC	ACAGTGCCTA 1320
1321	TGGGGAGCCC	AGGAAGCTGC	TCACCCAAGA	TTTGGTGCG	GAAAAGTACC	TGGAGTACCG 1380
1381	GCAGGTGCCG	GACAGTGATC	CCGCACGCTA	TGAGTTCCTG	TGGGGTCCAA	GGGCCCTCGC 1440
1441	TGAACCAGC	TATGTGAAAG	TCCTTGAGTA	TGTGATCAAG	GTCAGTGCAA	GAGTTCGCTT 1500
1501	TTTCTTCCCA	TCCCTGCCGTG	AAGCAGCTTT	GAGAGAGGAG	GAAGAGGGAG	TCTGAGCATG 1560
1561	AGTTGCAGCC	AAGGCCAGTG	GGAGGGGGAC	TGGGCCAGTG	CACCTTCCAG	GGCCCGGTCC 1620
1621	AGCAGCTTCC	CCTGCCCTCGT	GTGACATGAG	GCCCATTCTT	CACTCTGAAG	AGAGCGGTCA 1680
1681	GTGTCTCAG	TAGTAGGTTT	CTGTTCTATT	GGGTGACTTG	GAGATTTATC	TTTGTCTCTT 1740
1741	TTTGGAATTG	TTCAAATGTT	TTTTTTTAAG	GGATGGTTGA	ATGAACTTCA	GCATCCAAGT 1800
1801	TTATGAATGA	CAGCAGTCAC	ACAGTCTCTG	GTATATAGTT	TAAGGGTAAG	AGTCTTGTGT 1860
1861	TTTATTCAGA	TTGGGAAATC	CATTCTATTT	TGTGAATTGG	GATAATAACA	GCAGTGGAAAT 1920
1921	AAGTACTTAG	AAATGTGAAA	AATGAGCAGT	AAAATAGATG	AGATAAAGAA	CTAAAGAAAT 1980
1981	TAAGAGATAG	TCAATTCCTG	CCTTATACCT	CAGTCTATTC	TGTAATAATTT	TTAAAGATAT 2040
2041	ATGCATACCT	GGATTTCCTT	GGCTTCTTTG	AGAATGTAAG	AGAAATTAAA	TCTGAATAAA 2100
2101	GAATTCTTCC	TGTTCACTGG	CTCTTTTCTT	CTCCATGCAC	TGAGCATCTG	CTTTTGGAA 2160
2161	GGCCCTGGGT	TAGTAGTGGA	GATGCTAAGG	TAAGCCAGAC	TCATACCCAC	CCATAGGGTC 2220
2221	GTAGAGTCTA	GGAGCTGCAG	TCACGTAATC	GAGGTGGCAA	GATGTCCTCT	AAAGATGTAG 2280
2281	GGAAAAGTGA	GAGAGGGGTG	AGGGTGTGGG	GCTCCGGGTG	AGAGTGGTGG	AGTGTCAATG 2340
2341	CCCTGAGCTG	GGGCATTTTG	GGCTTTGGGA	AACTGCAGTT	CCTTCTGGGG	GAGCTGATTG 2400
2401	TAATGATCTT	GGGTGGATCC				2420
	10	20	30	40	50	60

**Example 21**

After the 2.4 kb genomic segment had been identified, studies were carried out to determine if an "E+" subline expressed any homologous DNA. Cell line MZ2-MEL 3.0 was used as a source, and a cDNA library was prepared from its mRNA, using art known techniques. The 2.4 kb segment was used as a probe, and mRNA of about 1.8 kb was identified as homologous, using Northern blot analysis. When cDNA was screened, clones were obtained showing almost complete identity to parts of the 2.4 kb fragment. Two exons were thus identified. An additional exon was located upstream of these, via sequencing segments of cosmid B3 located in front of the 2.4 kb BamHI fragment. The gene extends over about 4.5 kb, as shown in Figure 8. The starting point of the transcribed region was confirmed using PCR for the 5' end of the cDNA. The three exons comprise 65, 73, and 1551 base pairs. An ATG is located at position 66 of exon 3, followed by an 828 base pair reading frame.

**Example 22**

To determine if smaller segments of the 2.4 kb fragment could transfer the expression of antigen E, smaller pieces corresponding to the larger gene were prepared, using art recognized techniques, and transferred into E<sup>-</sup> cells. Figure 8 shows the boundaries of the three segments.

Transfer of antigen expression in this manner indicates that the gene codes for the antigen precursor, rather than coding for a protein which activates the antigen.

**Example 23**

The probing of cDNA described supra revealed, surprisingly, two different but closely related cDNAs. These cDNAs, when tested, did not transfer expression of antigen E, but they do show substantial homology to the first cDNA segment. The three segments, appear to indicate a newly recognized family of genes, referred to as "MAGE" for "melanoma antigen". In Figure 9, "mage-1" directs expression of the antigen from MZ2 cells. Portions of the third exon of each gene are presented in Figure 9. The second and third

sequences are more closely related to each other than the first (18.1 and 18.9% difference compared to the first; 12% with each other). Out of 9 cDNA clones obtained, three of each type were obtained, suggesting equal expression. "MAGE" as used hereafter refers to a family of molecules, and the nucleic acids coding for them. These nucleic acids share a certain degree of homology and are expressed in tumor cells including several types of human tumor cells as well as in human tumors. The family is referred to as "MAGE" because the first members were identified in human melanoma cells. As the experiments which follow indicate, however, the members of the MAGE family are not at all restricted to melanoma tumors; rather, MAGE refers to a family of tumor rejection antigen precursors and the nucleic acid sequences coding therefore. The antigens resulting therefrom are referred to herein as "MAGE TRAs" or "melanoma antigen tumor rejection antigens"

#### Example 24

Experiments with mouse tumors have demonstrated that new antigens recognized by T cells can result from point mutations that modify active genes in a region that codes for the new antigenic peptide. New antigens can also arise from the activation of genes that are not expressed in most normal cells. To clarify this issue for antigen MZ2-E, the mage-1 gene present in the melanoma cells was compared to that present in normal cells of patient MZ2.

Amplification by polymerase chain reaction (PCR) of DNA of phytohemagglutinin-activated blood lymphocytes using primers surrounding a 1300 bp stretch covering the first half of the 2.4 kb fragment was carried out. As expected, a PCR product was obtained whereas none was obtained with the DNA of the E<sup>-</sup> variant. The sequence of this PCR product proved identical to the corresponding sequence of the gene carried by the E<sup>+</sup> melanoma cells. Moreover, it was found that antigen MZ2-E was expressed by cells transfected with the cloned PCR product. This result suggests that the activation of a gene normally silent is responsible for the appearance of tumor rejection antigen MZ2-E.



Example 25

In order to evaluate the expression of gene mage-1 by various normal and tumor cells, Northern blots were hybridized with a probe covering most of the third exon. In contrast with the result observed with human tumor cell line MZ2-MEL 3.0, no band was observed with RNA isolated from a CTL clone of patient MZ2 and phytohemagglutinin-activated blood lymphocytes of the same patient. Also negative were several normal tissues of other individuals (Figure 10 and Figure 11). Fourteen melanoma cell lines of other patients were tested. Eleven were positive with bands of varying intensities. In addition to these culture cell lines, four samples of melanoma tumor tissue were analyzed. Two samples, including a metastasis of patient MZ2 proved positive, excluding the possibility that expression of the gene represented a tissue culture artefact. A few tumors of other histological types, including lung tumors were tested. Most of these tumors were positive (Figures 10 and 11). These results indicated that the MAGE gene family is expressed by many melanomas and also by other tumors. However, they provided no clear indication as to which of genes mage-1, 2 or 3 were expressed by these cells, because the DNA probes corresponding to the three genes cross-hybridized to a considerable extent. To render this analysis more specific, PCR amplification and hybridization with highly specific oligo- nucleotide probes were used. cDNAs were obtained and amplified by PCR using oligonucleotide primers corresponding to sequences of exon 3 that were identical for the three MAGE genes discussed herein. The PCR products were then tested for their ability to hybridize to three other oligonucleotides that showed complete specificity for one of the three genes (Figure 9). Control experiments carried out by diluting RNA of melanoma MZ2-MEL 3.0 in RNA from negative cells indicated that under the conditions used herein the intensity of the signal decreased proportionally to the dilution and that positive signals could still be detected at a dilution of 1/300. The normal cells (lymphocytes) that were tested by PCR were confirmed to be negative for the

expression of the three MAGE genes, suggesting therefore a level of expression of less than 1/300<sup>th</sup> that of the MZ2 melanoma cell line (Figure 11). For the panel of melanoma cell lines, the results clearly showed that some melanomas expressed MAGE genes mage 1, 2 and 3 whereas other expressed only mage-2 and 3 (Figures 11 and 10). Some of the other tumors also expressed all three genes whereas others expressed only mage-2 and 3 or only mage-3. It is impossible to exclude formally that some positive PCR results do not reflect the expression of one of the three characterized MAGE genes but that of yet another closely related gene that would share the sequence of the priming and hybridizing oligonucleotides. It can be concluded that the MAGE gene family is expressed by a large array of different tumors and that these genes are silent in the normal cells tested to this point.

#### Example 26

The availability of a sequence that transfects at high efficiency and efficiently expresses a TRAP made it possible to search for the associated major histocompatibility complex (MHC) class I molecule. The class I specificities of patient MZ2 are HLA-A1, A29, B37, B44 and C6. Four other melanomas of patients that had A1 in common with MZ2 were cotransfected with the 2.4 kb fragment and pSVtkneo $\beta$ . Three of them yielded neo<sup>r</sup> transfectants that stimulated TNF release by anti-E CTL clone 82/30, which is CD8<sup>+</sup> (Figure 10). No E- transfectant was obtained with four other melanomas, some of which shared A29, B44 or C6 with MZ2. This suggests that the presenting molecule for antigen MZ2-E is HLA-A1. In confirmation, it was found that, out of 6 melanoma cell lines derived from tumors of HLA-A1 patients, two stimulated TNF release by anti-E CTL clone 82/30 of patient MZ2. One of these tumor cell lines, MI13443-MEL also showed high sensitivity to lysis by these anti-E CTL. These two melanomas were those that expressed mage-1 gene (Figure 13). Eight melanomas of patients with HLA haplotypes that did not include A1 were examined for their sensitivity to lysis and for their ability to stimulate TNF release by the CTL. None was found to be positive. The

ability of some human anti-tumor CTL to lyse allogeneic tumors sharing an appropriate HLA specificity with the original tumor has been reported previously (Darrow, et al., J. Immunol. 142: 3329 (1989)). It is quite possible that antigenic peptides encoded by genes mage 2 and 3 can also be presented to autologous CTL by HLA-A1 or other class I molecules, especially in view of the similar results found with murine tumors, as elaborated upon supra.

#### Example 27

As indicated supra, melanoma MZ2 expressed antigens F, D and A', in addition to antigen E. Following the isolation of the nucleic acid sequence coding for antigen E, similar experiments were carried out to isolate the nucleic acid sequence coding for antigen F.

To do this, cultures of cell line MZ2-MEL2.2, an E<sup>-</sup> cell line described supra, were treated with anti-F CTL clone 76/6, in the same manner described for treatment with anti-E CTL clones. This resulted in the isolation of an F antigen loss variant, which was then subjected to several rounds of selection. The resulting cell line, "MZ2-MEL2.2.5" was completely resistant to lysis by anti-F CTLs, yet proved to be lysed by anti-D CTLs.

Again, following the protocols set forth for isolation of antigen -E precursor DNA, the F<sup>-</sup> variant was transfected with genomic DNA from F<sup>+</sup> cell line MZ2-MEL3.0. The experiments yielded 90,000 drug resistant transfectants. These were tested for MZ2-F expression by using pools of 30 cells in the TNF detection assay elaborated upon supra. One pool stimulated TNF release by anti-F CTLs, and was cloned. Five of 145 clones were found to stimulate anti-F CTLs. Lysis assays, also following protocols described supra, confirmed (i) expression of the gene coding for antigen F, and (ii) presentation of antigen F itself.

#### Example 28

Following identification of F<sup>+</sup> cell lines, the DNA therefrom was used to transfect cells. To do this, a cosmid library of F<sup>+</sup> cell line MZ2-MEL.43 was prepared, again using

the protocols described supra. The library was divided into 14 groups of about 50,000 cosmids, and DNA from each group was transfected into MZ2-MEL2.2.5. Transfectants were then tested for their ability to stimulate TNF release from anti-F CTL clone 76/6. Of 14 groups of cosmids, one produced two independent transfectants expressing antigen F; a yield of two positives out of 17,500 genitacin resistant transfectants.

#### Example 29

The existence of a gene family was suggested by the pattern observed on the Southern blot (Figure 12). To do this, the 2.4 kb BamHI fragment, which transferred the expression of antigen M22-E, was labelled with 32p and used as a probe on a Southern Blot of BamHI digested DNA of E + cloned subclone M22-MEL2.2. Hybridization conditions included 50  $\mu$ l/cm<sup>2</sup> of 3.5xSSC, 1xDenhardt's solution; 25 mM sodium phosphate buffer (pH 7.0), 0.5% SDS, 2mM EDTA, where the 2.4 kb probes had been labelled with [ $\alpha^{32}$ p]dCTP (2-3000 Ci/mole), at  $3 \times 10^6$  cpm/ml. Hybridization was carried out for 18 hours at 65°C. After this, the membranes were washed at 65°C four times for one hour each in 2xSSC, 0.1% SDS, and finally for 30 minutes in 0.1xSSC, 0.1% SDS. To identify hybridization, membranes were autoradiographed using Kodak X-AR film and Kodak X-Omatic fine intensifying screens.

In the following examples, whenever "hybridization" is referred to, the stringency conditions used were similar to those described supra. "Stringent conditions" as used herein thus refers to the foregoing conditions; subject to routine, art recognized modification.

#### Example 30

The cDNA coding for mage 4 was identified from a sample of the human sarcoma cell line LB23-SAR. This cell line was found to not express mage 1, 2 or 3, but the mRNA of the cell line did hybridize to the 2.4 kb sequence for mage 1. To study this further, a cDNA library was prepared from total LB23-SAR mRNA, and was then hybridized to the 2.4 kb fragment. A cDNA sequence was identified as hybridizing to this probe, and is identified hereafter as mage 4.

Example 31

Experiments were carried out using PHA-activated lymphocytes from patient "MZ2", the source of the "MZ" cells discussed supra. An oligonucleotide probe which showed  
5 homology to mage 1 but not mage 2 or 3 was hybridized with a cosmid library derived from the PHA activated cells. The size of the hybridizing BamHI cosmid fragment, however, was 4.5 kb, thus indicating that the material was not mage 1; however, on the basis of homology to mage 1-4, the fragment can be  
10 referred to as "mage 5". The sequence of MAGE 5 is presented in SEQ ID NO: 16.

Example 32

Melanoma cell line LB-33-MEL was tested. Total mRNA from the cell line was used to prepare cDNA, which was then  
15 amplified with oligos CHO9: (ACTCAGCTCCTCCCAGATTT), and CHO10: (GAAGAGGAGGGGCCAAG). These oligos correspond to regions of exon 3 that are common to previously described mage 1, 2 and 3.

To do this, 1  $\mu$ g of RNA was diluted to a total volume of  
20 20  $\mu$ l, using 2  $\mu$ l of 10x PCR buffer, 2  $\mu$ l of each of 10 mM dNTP, 1.2  $\mu$ l of 25 mM  $MgCl_2$ , 1  $\mu$ l of an 80 mM solution of CHO9, described supra, 20 units of RNasin, and 200 units of M-MLV reverse transcriptase. This was followed by incubation for 40 minutes at 42°C. PCR amplification followed, using 8  
25  $\mu$ l of 10x PCR buffer, 4.8  $\mu$ l of 25 mM  $MgCl_2$ , 1  $\mu$ l of CHO10, 2.5 units of *Thermus aquaticus* ("Taq") polymerase, and water to a total volume of 100  $\mu$ l. Amplification was then carried out for 30 cycles (1 minute 94°C; 2 minutes at 52°C, 3 minutes at 72°C). Ten  $\mu$ l of each reaction were then size fractionated  
30 on agarose gel, followed by nitrocellulose blotting. The product was found to hybridize with oligonucleotide probe CHO18 (TCTTGTATCCTGGAGTCC). This probe identified mage 1 but not mage 2 or 3. However, the product did not hybridize to probe SEQ 4 (TTGCCAAGATCTCAGGAA). This probe also binds mage  
35 1 but not 2 and 3. This indicated that the PCR product contained a sequence that differed from mage 1, 2 and 3. Sequencing of this fragment also indicated differences with

respect to mage 4 and 5. These results indicate a sequence differing from previously identified mage 1, 2, 3, 4 and 5, and is named mage 6.

Example 33

5 In additional experiments using cosmid libraries from PHA-activated lymphocytes of MZ2, the 2.4 kb mage 1 fragment was used as a probe and isolated a complementary fragment. This clone, however, did not bind to oligonucleotides specific for mage 1, 2, 3 or 4. The sequence obtained shows some  
10 homology to exon 3 of mage 1, and differs from mages 1-6. It is referred to as mage 7 hereafter. Additional screenings yielded mage 8-11.

Example 34

15 The usefulness of the TRAPs, as well as TRAs derived therefrom, was exemplified by the following.

Exon 3 of mage 1 was shown to transfer expression of antigen E. As a result, it was decided to test whether synthetic peptides derived from this exon 3 could be used to confer sensitivity to anti-E CTL.

20 To do this, and using standard protocols, cells normally insensitive to anti-E/CTLs were incubated with the synthetic peptides derived from Exon 3.1. Using the CTL lytic assays described supra on P815A, and a peptide concentration of 3 mM, the peptide Glu-Ala-Asp-Pro-Thr-Gly-His-Ser-Tyr was shown to  
25 be best. The assay showed lysis of 30%, indicating conferring of sensitivity to the anti-E CTL.

Example 35

Nucleic acid sequences referred to as "smage" were isolated from murine cells. Using the protocols described  
30 supra, a cosmid library was prepared from the DNA of normal DBA/2 kidney cells, using cosmid vector C2RB. As a probe, the 2.4 kb BamHI fragment of MAGE-1 was used. The DNA was blotted to nylon filters, and these were washed in 2xSSC at 65°C to identify the smage material.

35 Example 36

Further tissue samples were tested for the presence of MAGE genes, using the protocols discussed supra. Some of

these results follow.

There was no expression of the MAGE genes in brain or kidney tumor tissue. Colon tumor tissue showed expression of MAGE 1, 2, 3 and 4, although not all tumors tested showed expression of all MAGE genes. This is also true for pancreatic tumor (MAGE 1); non-small cell lung (MAGE 1, 2, 3 and 4), prostate (MAGE 1), sarcomas (MAGE 1, 2, 3 and 4), breast (MAGE 1, 2 and 3), and larynx (MAGE 1 and 4).

#### Example 37

A cytolytic CTL clone "20/38" was obtained from peripheral blood lymphocytes of melanoma patient MZ2. This clone is described by Van den Eynde et al., Int. J. Cancer 44: 634-640 (1989), the disclosure of which is incorporated by reference. The CTL clone has isolated following Herin et al., Int. J. Cancer 39: 390-396 (1987), which is incorporated by reference. The assay is described herein, however. Autologous melanoma cells were grown in vitro, and then resuspended at  $10^7$  cells/ml in DMEM, supplemented with 10% HEPES and 30% FCS, and incubated for 45 minutes at 37°C with 200  $\mu$ Ci/ml of  $\text{Na}^{(51)\text{Cr}}\text{O}_4$ . Labelled cells were washed three times with DMEM, supplemented with 10 mM HEPES. These were then resuspended in DMEM supplemented with 10 mM HEPES and 10% FCS, after which 100  $\mu$ l aliquots containing  $10^3$  cells, were distributed into 96 well microplates. Samples of the CTL clone were added in 100  $\mu$ l of the same medium, and assays were carried out in duplicate. Plates were centrifuged for four minutes at 100g, and incubated for four hours at 37°C in a 5.5%  $\text{CO}_2$  atmosphere.

Plates were centrifuged again, and 100  $\mu$ l aliquots of supernatant were collected and counted. Percentage of  $^{51}\text{Cr}$  release was calculated as follows:

$$\% \text{ } ^{51}\text{Cr} \text{ release} = \frac{(\text{ER}-\text{SR})}{(\text{MR}-\text{SR})} \times 100$$

where ER is observed, experimental  $^{51}\text{Cr}$  release, SR is spontaneous release measured by incubating  $10^3$  labeled cells in 200  $\mu$ l of medium alone, and MR is maximum release, obtained

by adding 100 ul 0.3% Triton X-100 to target cells.

Those mononuclear blood samples which showed high CTL activity were expanded and cloned via limiting dilution, and were screened again, using the same methodology.

5       The same method was used to test target K562 cells. When EBV-B cells were used, the only change was the replacement of DMEM medium by Hank's medium, supplemented with 5% FCS.

These experiments led to isolation of CTL clone 20/38.

10       Figure 1 presents the results of these assays. Specifically, it will be seen that the CTL clone lysed autologous melanoma cell line MZ2-MEL.3.0, but did not lyse EBV-B cell lines, fibroblasts, K562 or non-autologous melanoma cell line SK-MEL-29.

15       Example 38

Once the CTL clone was recognized as being specific for the autologous cell line, it was tested for antigenic specificity. To do this, antigen loss variants derived from patient MZ2 were tested in the same type of chromium release assay described above. These target lines were MZ2-MEL 3.0, which is D<sup>+</sup>, E<sup>+</sup>, F<sup>+</sup>, A<sup>+</sup>, MZ2-MEL.61, which is D<sup>-</sup>, MZ2-MEL 2.2, which is E<sup>-</sup>, and MZ2-MEL.4, which is F<sup>-</sup>. In addition to CTL clone 20/38, clones which are known to be anti-A (CTL 28/336), anti-F (CTL 76/6), and anti-E (CTL 22/13) were tested.

25       These results are set forth in figure 15. It will be seen that CTL clone 20/38 lysed all the cell lines leading to chromium release except D<sup>-</sup> cell line MZ2-MEL.61, thus indicating that the CTL clone is anti-D. This result was confirmed, in experiments not included herein, by experiments where TNF release by the CTL clone was observed only in the presence of melanoma lines presenting antigen D.

30       Example 39

Once antigen D was identified as the target molecule, studies were carried out to determine the HLA type which presented it. The experiments described in example A showed that antigen D was presented by MZ2-MEL, and this cell line's HLA specificity is known (i.e., A1, A29, B37, B44, Cw6,

35



C.cl.10). It was also known, however, that a variant of MZ2-MEL which had lost HLA molecules A29, B44 and C.cl.10 still expressed antigen D, so these could be eliminated from consideration. Studies were not carried out on lines  
5 expressing B37, as none could be found.

In all, 13 allogeneic lines were tested, which expressed either HLA-A1 (10 of 13), or Cw6 (3 of 13). The cell lines were tested for their ability to stimulate release of TNF by CTL clone 20/38, using the method of Traversari et al.,  
10 Immunogenetics 35: 145-152 (1992), the disclosure of which is incorporated by reference. This assay measures TNF release via testing toxicity of supernatants on WEHI 164-13 cells.

In the assays, cell samples (3000, 10,000 or 30,000 cells) from the allogeneic lines were cultured in the presence  
15 of 1500 cells of the CTL clone, and 25 u/ml of IL-2. Twenty-four hours later, the supernatant from the culture was tested against the WEHI cells for toxicity. The results are presented in Table 3, which follows.

Eight cell lines were found to stimulate TNF release from the CTL clone 20/38. All of these lines were HLA-A1. None of  
20 the Cw6 presenting lines did so.

The cell lines were also assayed to determine MAGE expression. All eight of the lines which stimulated TNF release expressed MAGE-3, whereas the two HLA-A1 lines which  
25 were negative did not.

Table 3

Melanoma	Number of Cells	TNF pg/ml		Expression of Mage-3	Expression of HLA-A-1
		Exp 1		Exp 2	
		+CTL 20/38		+CTL 20/38	
MZ2-MEL.61.2	50000	1		4	+++ +
MZ2-MEL-ET1	50000	>120		>120	+++ +
	1666	66		>120	
LY-1-MEL	30000	1	>120	1	+++ +
	10000	1	>120	1	
	3000	<1	114	2	
MI-10221	30000	<1	>120		+++ +
	10000	<1	71		
	3000	<1	74		
LY-2-MEL	30000	1	57		+++ +
	10000	1	86		
	3000	1	91		
LY-4-MEL	30000	1	>120		+++ +
	10000	1	>120		
	3000	1	>120		
SK23-MEL	30000	1	112		+++ +
	10000	1	116		
	3000	1	105		
MI-665/2-MEL	30000	1	3	2 4	- +
	10000	1	2	2 5	
	3000	1	5,2	1 5	
LB34-MEL	30000	1	>120		+++ +
	10000	1	>120		
	3000	1	>120		
LB45-MEL	30000	1	11	1 30	- +
	10000	1	6	1 12	
	3000	1	2	<1 7	
NA-6-MEL	30000	1	77	5 98	+++ +
	10000	1	104	5 >120	
	3000	1	110	4 >120	
MI-13443-MEL	30000	1	>120		+++ +
	10000	1	>120		
	3000	1	>120		
LB5-MEL	30000	1	8	4 9	+ -
	10000	<1	5	4 11	
	3000	<1	5	1 5	
SK64-MEL	30000	1	4	2 5	? -
	10000	1	2	1 5	
	3000	1	1	1 4	
LB33-MEL	30000			1 3,5	+++ -
	10000			1 4	
	3000			1 3	
LB73-MEL	50000		16		- -

1500 CTL 20/38 and 25 $\mu$ /ml IL2 were mixed with the indicated number of cells of the different allogeneic melanomas. 24 hours later, the amount of TNF present in the supernatant was assayed by testing its cytotoxicity for WEHI-164-13 cells.

SUBSTITUTE SHEET (RULE 26)

Example 40

In view of the results set forth in example C, experiments were carried out to determine if antigen D was in fact a tumor rejection antigen derived from MAGE-3. To do this, recipient COS7 cells were transfected with 100ng of the gene for HLA-A1 cloned into pcDNA I/Amp, and 100 ng of one of (a) cDNA for MAGE-1 cloned into pcDNA I/Amp, (b) cDNA for MAGE-2 cloned into pcDSR $\alpha$ , or (c) cDNA for MAGE-3 cloned into pcDSR $\alpha$ . The transfecting sequences were ligated into the plasmids in accordance with manufacturer's instructions. Samples of COS-7 cells were seeded, at 15,000 cells/well into tissue culture flat bottom microwells, in Dulbecco's modified Eagles Medium ("DMEM") supplemented with 10% fetal calf serum. The cells were incubated overnight at 37°C, medium was removed and then replaced by 30  $\mu$ l/well of DMEM medium containing 10% Nu serum, 400  $\mu$ g/ml DEAE-dextran, 100  $\mu$ M chloroquine, and the plasmids described above. Following four hours of incubation at 37°C, the medium was removed, and replaced by 50  $\mu$ l of PBS containing 10% DMSO. This medium was removed after two minutes and replaced by 200  $\mu$ l of DMEM supplemented with 10% of FCS.

Following this change in medium, COS cells were incubated for 24 hours at 37°C. Medium was then discarded, and 1500 cells of CTL clones 20/38 were added, in 100  $\mu$ l of Iscove medium containing 10% pooled human serum, supplemented with 25 u/ml of IL-2. Supernatant was removed after 24 hours, and TNF content was determined in an assay on WEHI cells, as described by Traversari et al., Immunogenetics 35: 145-152 (1992), the disclosure of which is incorporated by reference. These results are shown in Figure 16.

It will be seen that the CTL clone was strongly stimulated by COS7 cells transfected with HLA-A1 and MAGE-3, but not by the cells transfected with the other mage genes. This leads to the conclusion that antigen D is a tumor rejection antigen derived from the tumor rejection antigen precursor coded by gene MAGE-3, and that this TRA is presented

by HLA-A1 molecules.

The foregoing disclosure, including the examples, places many tools of extreme value in the hands of the skilled artisan. To begin, the examples identify and provide a methodology for isolating nucleic acid molecules which code for tumor rejection antigen precursors as well as the nucleic acid molecules complementary thereto. It is known that DNA exists in double stranded form, and that each of the two strands is complementary to the other. Nucleic acid hybridization technology has developed to the point where, given a strand of DNA, the skilled artisan can isolate its complement, or synthesize it.

"Nucleic acid molecule" as used herein refers to all species of DNA and RNA which possess the properties discussed supra. Genomic and complementary DNA, or "cDNA" both code for particular proteins, and as the examples directed to isolation of MAGE coding sequences show, this disclosure teaches the artisan how to secure both of these.

Similarly, RNA molecules, such as mRNA can be secured. Again, with reference to the skilled artisan, once one has a coding sequence in hand, mRNA can be isolated or synthesized.

Complementary sequences which do not code for TRAP, such as "antisense DNA" or mRNA are useful, e.g., in probing for the coding sequence as well as in methodologies for blocking its expression.

It will also be clear that the examples show the manufacture of biologically pure cultures of cell lines which have been transfected with nucleic acid sequences which code for or express the TRAP molecules. Such cultures can be used as a source for tumor rejection antigens, e.g., or as therapeutics. This aspect of the invention is discussed infra.

Cells transfected with the TRAP coding sequences may also be transfected with other coding sequences. Examples of other coding sequences include cytokine genes, such as interleukins (e.g., IL-2 or IL-4), or major histocompatibility complex (MHC) or human leukocyte antigen (HLA) molecules.

Cytokine gene transfection is of value because expression of these is expected to enhance the therapeutic efficacy of the biologically pure culture of the cells in vivo. The art is well aware of therapies where interleukin transfectants have been administered to subjects for treating cancerous conditions. In a particularly preferred embodiment, cells are transfected with sequences coding for each of (i) a TRAP molecule, (ii) an HLA/MHC molecule, and (iii) a cytokine.

Transfection with an MHC/HLA coding sequence is desirable because certain of the TRAs may be preferentially or specifically presented only by particular MHC/HLA molecules. Thus, where a recipient cell already expresses the MHC/HLA molecule associated with presentation of a TRA, additional transfection may not be necessary although further transformation could be used to cause over-expression of the antigen. On the other hand, it may be desirable to transfect with a second sequence when the recipient cell does not normally express the relevant MHC/HLA molecule. It is to be understood, of course, that transfection with one additional sequence does not preclude further transfection with other sequences.

The term "biologically pure" as used in connection with the cell line described herein simply means that these are essentially free of other cells. Strictly speaking, a "cell line" by definition is "biologically pure", but the recitation will establish this fully.

Transfection of cells requires that an appropriate vector be used. Thus, the invention encompasses expression vectors where a coding sequence for the TRAP of interest is operably linked to a promoter. The promoter may be a strong promoter, such as those well known to the art, or a differential promoter, i.e., one which is operative only in specific cell types. The expression vectors may also contain all or a part of a viral or bacterial genome, such as vaccinia virus or BCG. Such vectors are especially useful in preparing vaccines.

The expression vectors may incorporate several coding sequences, as long as the TRAP sequence is contained therein.

The cytokine and/or MHC/HLA genes discussed supra may be included in a single vector with the TRAP sequence. Where this is not desired, then an expression system may be provided, where two or more separate vectors are used where each coding sequence is operably linked to a promoter. Again, the promoter may be a strong or differential promoter. Co-transfection is a well known technique, and the artisan in this field is expected to have this technology available for utilization. The vectors may be constructed so that they code for the TRA molecule directly, rather than the TRAP molecule. This eliminates the need for post-translational processing.

As the foregoing discussion makes clear, the sequences code for "tumor rejection antigen precursors" ("TRAPs") which, in turn, are processed into tumor rejection antigens ("TRAs"). Isolated forms of both of these categories are described herein, including specific examples of each. Perhaps their most noteworthy aspect is as vaccines for treating various cancerous conditions. The evidence points to presentation of TRAs on tumor cells, followed by the development of an immune response and deletion of the cells. The examples show that when various TRAs are administered to cells, a CTL response is mounted and presenting cells are deleted. This is behavior characteristic of vaccines, and hence TRAPs, which are processed into TRAs, and the TRAs themselves may be used, either alone or in pharmaceutically appropriate compositions, as vaccines. Similarly, presenting cells may be used in the same manner, either alone or as combined with ingredients to yield pharmaceutical compositions. Additional materials which may be used as vaccines include isolated cells which present the TRA molecule on their surface, as well as TRAP fragments, mutated viruses, especially etioloated forms, and transfected bacteria. "Fragments" as used herein refers to peptides which are smaller than the TRA, but which possess the properties required of a vaccine, as discussed supra. Another vaccine comprises or consists of complexes of TRA and HLA molecule. Vaccines of the type described herein may be used preventively, i.e., via administration to a subject in an

amount sufficient to prevent onset of a cancerous condition.

The generation of an immune response, be it T-cell or B-cell related, is characteristic of the effect of the presented tumor rejection antigen. With respect to the B-cell response, this involves, inter alia, the generation of antibodies to the TRA, i.e., which specifically bind thereto. In addition, the TRAP molecules are of sufficient size to render them immunogenic, and antibodies which specifically bind thereto are a part of this invention. These antibodies may be polyclonal or monoclonal, the latter being prepared by any of the well recognized methodologies for their preparation which need not be repeated here. For example, mAbs may be prepared using an animal model, e.g., a Balb/C mouse or in a test tube, using, e.g., EBV transformants. In addition, antiserum may be isolated from a subject afflicted with a cancerous condition where certain cells present a TRA. Such antibodies may also be generated to epitopes defined by the interaction of TRA and HLA/MHC molecules.

Review of the foregoing disclosure will show that there are a number of facets to the system which may be referred to as "tumor rejection antigen presentation and recognition". Recognition of these phenomena has diagnostic consequences. For example, the existence of specific CTL clones, or antibodies to the TRA makes it possible to diagnose or monitor cancerous conditions (explained infra), by monitoring the CTLs in a sample from a subject, binding of antibodies to TRAs, or the activity of anti-TRA CTLs in connection with subject samples. Similarly, the expression of nucleic acid molecules for TRAPs can be monitored via amplification (e.g., "polymerase chain reaction"), anti-sense hybridization, probe technologies, and so forth. Various subject samples, including body fluids (blood, serum, and other exudates, e.g.), tissues and tumors may be so assayed.

A particular manner of diagnosis is to use an adaptation of the standard "tuberculin test" currently used for diagnosis of tuberculosis. This standard skin test administers a stable form of "purified protein derivative" or "PPD" as a diagnostic

aid. In a parallel fashion, TRAs in accordance with this invention may be used in such a skin test as a diagnostic aid or monitoring method.

5 The term "cancerous condition" is used herein to embrace all physiological events that commence with the initiation of the cancer and result in final clinical manifestation. Tumors do not spring up "ab initio" as visible tumors; rather there are various events associated with the transformation of a normal cell to malignancy, followed by development of a growth  
10 of biomass, such as a tumor, metastasis, etc. In addition, remission may be conceived of as part of "a cancerous condition" as tumors seldom spontaneously disappear. The diagnostic aspects of this invention include all events involved in carcinogenesis, from the first transformation to malignancy of a single cell, through tumor development and  
15 metastasis, as well as remission. All are embraced herein.

Where "subject" is used, the term embraces any species which can be afflicted with a cancerous condition. This includes humans and non-humans, such as domesticated animals,  
20 breeding stock, and so forth.

There are therapeutic aspects of this invention as well. The efficacy of administration of effective amounts of TRAPs and TRAs as vaccines has already been discussed supra. Similarly, one may develop the specific CTLs in vitro and then  
25 administer these to the subject. Antibodies may be administered, either polyclonal or monoclonal, which specifically bind to cells presenting the TRA of interest. These antibodies may be coupled to specific antitumor agents, including, but not being limited to, methotrexate radio-  
30 iodinated compounds, toxins such as ricin, other cytostatic or cytolytic drugs, and so forth. Thus, "targeted" antibody therapy is included herein, as is the application of deletion of the cancerous cells by the use of CTLs.

The data from examples 37-40 show that a tumor rejection antigen derived from MAGE-3 is presented by HLA-A1 molecules.  
35 As such, in addition to the nucleic acid molecules coding for this TRAP, the TRAP itself as coded for by the sequences,



vectors, cell lines, etcetera which incorporate this nucleic acid molecule, the invention also encompasses combination of the molecules coding for the MAGE-3 TRAP and HLA-A1. Thus, co-transfectants, vectors containing coding sequences for both, expression systems such as kits, or separate vectors, and so forth, are all embraced by the invention. Similarly, the vaccines discussed supra can be made by incorporating the TRAP from MAGE-3 and an adjuvant.

It is to be understood that a given TRAP may yield more than one TRA. In the case of MAGE-3, it has been shown that antigen D, as the term is used herein, derives therefrom, and one aspect of the invention is this isolated tumor rejection antigen. Another is isolated complexes of the TRA and its presenting molecule, i.e., HLA-A1.

The identification of MAGE-3 derived TRAs as being presented by HLA-A1 molecules suggests various therapeutic and diagnostic approaches. In a therapeutic context, e.g., the treatment of a disorder characterized by MAGE-3 expression may be treated in a number of ways, "disorder" being used to refer to any pathological condition where MAGE-3 TRAP is expressed, such as cancer (e.g., melanoma).

Therapeutic approaches based upon the disclosure are premised on a response by a subject's immune system, leading to lysis of TRA presenting cells, such as HLA-A1 cells. One such approach is the administration of CTLs specific to the complex to a subject with abnormal cells of the phenotype at issue. It is within the skill of the artisan to develop such CTLs in vitro. Specifically, a sample of cells, such as blood cells, are contacted to a cell presenting the complex and capable of provoking a specific CTL to proliferate. The target cell can be a transfectant, such as a COS cell of the type described supra. These transfectants present the desired complex on their surface and, when combined with a CTL of interest, stimulate its proliferation. COS cells, such as those used herein are widely available, as are other suitable host cells.

To detail the therapeutic methodology, referred to as

adoptive transfer (Greenberg, J. Immunol. 136(5): 1917 (1986); Reddel et al., Science 257: 238 (7-10-92); Lynch et al., Eur. J. Immunol. 21: 1403-1410 (1991); Kast et al., Cell 59: 603-614 (11-17-89)), cells presenting the desired complex are combined with CTLs leading to proliferation of the CTLs specific thereto. The proliferated CTLs are then administered to a subject with a cellular abnormality which is characterized by certain of the abnormal cells presenting the particular complex. The CTLs then lyse the abnormal cells, thereby achieving the desired therapeutic goal.

The foregoing therapy assumes that at least some of the subject's abnormal cells present the HLA/TRA complex. This can be determined very easily, as the art is very familiar with methods for identifying cells which present a particular HLA molecule, as well as how to identify cells expressing DNA containing the indicated sequences. Once isolated, such cells can be used with a sample of a subject's abnormal cells to determine lysis in vitro. If lysis is observed, then the use of specific CTLs in such a therapy may alleviate the condition associated with the abnormal cells. A less involved methodology examines the abnormal cells for HLA phenotyping, using standard assays, and determines expression via amplification using, e.g., PCR.

Adoptive transfer is not the only form of therapy that is available in accordance with the invention. CTLs can also be provoked in vivo, using a number of approaches. One approach, i.e., the use of non-proliferative cells expressing the complex, has been elaborated upon supra. The cells used in this approach may be those that normally express the complex, such as irradiated melanoma cells or cells transfected with one or both of the genes necessary for presentation of the complex. Chen et al., Proc. Natl. Acad. Sci. USA 88: 110-114 (January, 1991) exemplifies this approach, showing the use of transfected cells expressing HPVE7 peptides in a therapeutic regime. Various cell types may be used. Similarly, vectors carrying one or both of the genes of interest may be used. Viral or bacterial vectors are especially preferred. In these

systems, the gene of interest is carried by, e.g., a Vaccinia virus or the bacteria BCG, and the materials de facto "infect" host cells. The cells which result present the complex of interest, and are recognized by autologous CTLs, which then proliferate. A similar effect can be achieved by combining the tumor rejection antigen or the precursor itself with an adjuvant to facilitate incorporation into HLA-A1 presenting cells which present the HLA molecule of interest. The TRAP is processed to yield the peptide partner of the HLA molecule while the TRA is presented without the need for further processing. Thus, one may treat disorders where a MAGE-3 derived TRA is presented by HLA-A1 molecules, or by any HLA molecule.

In a diagnostic context, one may determine a disorder, as the term is used herein, by assaying for expression of the TRAP. This can be done directly (via, e.g., a PCR assay for TRAP sequences), or indirectly, via assaying for a MAGE-3 derived TRA, as the TRA's presence means that the TRAP is or was expressed.

It will be noted that two nucleic acid molecules are presented herein, i.e., MAGE-3 and MAGE-31, each of which code for TRAP MAGE-3. It is to be understood that when the expression "nucleic acid molecule which codes for MAGE-3 TRAP" is used, all molecules are covered which yield this molecule upon expression. Any number of variations, such as those showing codon degeneracy within the coding region, or variation within the introns, are covered by the invention.

The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, it being recognized that various modifications are possible within the scope of the invention.

## (1) GENERAL INFORMATION:

- (i) APPLICANTS: Gaugler, Béatrice; Van den Eynde, Benoît;  
van der Bruggen, Pierre; Boon-Falleur, Thierry
- (ii) TITLE OF INVENTION: Isolated Nucleic Acid Molecules Coding For  
Tumor Rejection Antigen Precursor Mage-3 And Uses Thereof
- (iii) NUMBER OF SEQUENCES: 26
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Felfe & Lynch
  - (B) STREET: 805 Third Avenue
  - (C) CITY: New York City
  - (D) STATE: New York
  - (F) ZIP: 10022
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Diskette, 5.25 inch, 360 kb storage
  - (B) COMPUTER: IBM
  - (C) OPERATING SYSTEM: PC-DOS
  - (D) SOFTWARE: Wordperfect
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: 08/037,230
  - (B) FILING DATE: 26-MARCH-1993
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: PCT/US92/04354
  - (B) FILING DATE: 22-MAY-1992
- (viii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 07/807,043
  - (B) FILING DATE: 12-DECEMBER-1991
- (ix) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 07/764,364
  - (B) FILING DATE: 23-SEPTEMBER-1991
- (x) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 07/728,838
  - (b) FILING DATE: 9-JULY-1991
- (xi) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 07/705,702
  - (B) FILING DATE: 23-MAY-1991
- (xii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Hanson, Norman D.
  - (B) REGISTRATION NUMBER: 30,946
  - (C) REFERENCE/DOCKET NUMBER: LUD 253.5
- (xiii) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (212) 688-9200
  - (B) TELEFAX: (212) 838-3884

- (2) INFORMATION FOR SEQUENCE ID NO: 1:  
(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 462 base pairs  
    (B) TYPE: nucleic acid  
    (D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: genomic DNA  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```
ACCACAGGAG AATGAAAAGA ACCCGGGACT CCCAAAGACG CTAGATGTGT GAAGATCCTG 60
ATCACTCATT GGGTGTCTGA GTTCTGCGAT ATTCATCCCT CAGCCAATGA GCTTACTGTT 120
CTCGTGGGGG GTTTGTGAGC CTTGGGTAGG AAGTTTTGCA AGTTCCGCCT ACAGCTCTAG 180
CTTGTGAATT TGTACCCTTT CACGTAAAAA AGTAGTCCAG AGTTTACTAC ACCCTCCCTC 240
CCCCCTCCCA CCTCGTGCTG TGCTGAGTTT AGAAGTCTTC CTTATAGAAG TCTTCCGTAT 300
AGAACTCTTC CGGAGGAAGG AGGGAGGACC CCCCCCTTT GCTCTCCCAG CATGCATTGT 360
GTCAACGCCA TTGCACTGAG CTGGTCGAAG AAGTAAGCCG CTAGCTTGCG ACTCTACTCT 420
TATCTTAACT TAGCTCGGCT TCCTGCTGGT ACCCTTTGTG CC 462
```

- (2) INFORMATION FOR SEQUENCE ID NO: 2:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 675 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

```

ATG TCT GAT AAC AAG AAA CCA GAC AAA GCC CAC AGT GGC TCA GGT GGT 48
Met Ser Asp Asn Lys Lys Pro Asp Lys Ala His Ser Gly Ser Gly Gly
      5      10      15
GAC GGT GAT GGG AAT AGG TGC AAT TTA TTG CAC CGG TAC TCC CTG GAA 96
Asp Gly Asp Gly Asn Arg Cys Asn Leu Leu His Arg Tyr Ser Leu Glu
      20      25      30
GAA ATT CTG CCT TAT CTA GGG TGG CTG GTC TTC GCT GTT GTC ACA ACA 144
Glu Ile Leu Pro Tyr Leu Gly Trp Leu Val Phe Ala Val Val Thr Thr
      35      40      45
AGT TTT CTG GCG CTC CAG ATG TTC ATA GAC GCC CTT TAT GAG GAG CAG 192
Ser Phe Leu Ala Leu Gln Met Phe Ile Asp Ala Leu Tyr Glu Glu Gln
      50      55      60
TAT GAA AGG GAT GTG GCC TGG ATA GCC AGG CAA AGC AAG CGC ATG TCC 240
Tyr Glu Arg Asp Val Ala Trp Ile Ala Arg Gln Ser Lys Arg Met Ser
      65      70      75
TCT GTC GAT GAG GAT GAA GAC GAT GAG GAT GAT GAG GAT GAC TAC TAC 288
Ser Val Asp Glu Asp Glu Asp Asp Glu Asp Asp Glu Asp Asp Tyr Tyr
      85      90      95
GAC GAC GAG GAC GAC GAC GAC GAT GCC TTC TAT GAT GAT GAG GAT GAT 336
Asp Asp Glu Asp Asp Asp Asp Asp Ala Phe Tyr Asp Asp Glu Asp Asp
      100      105      110
GAG GAA GAA GAA TTG GAG AAC CTG ATG GAT GAT GAA TCA GAA GAT GAG 384
Glu Glu Glu Glu Leu Glu Asn Leu Met Asp Asp Glu Ser Glu Asp Glu
      115      120      125
GCC GAA GAA GAG ATG AGC GTG GAA ATG GGT GCC GGA GCT GAG GAA ATG 432
Ala Glu Glu Glu Met Ser Val Glu Met Gly Ala Gly Ala Glu Glu Met
      130      135      140
GGT GCT GGC GCT AAC TGT GCC TGT GTT CCT GGC CAT CAT TTA AGG AAG 480
Gly Ala Gly Ala Asn Cys Ala Cys Val Pro Gly His His Leu Arg Lys
      145      150      155
AAT GAA GTG AAG TGT AGG ATG ATT TAT TTC TTC CAC GAC CCT AAT TTC 528
Asn Glu Val Lys Cys Arg Met Ile Tyr Phe Phe His Asp Pro Asn Phe
      165      170      175
CTG GTG TCT ATA CCA GTG AAC CCT AAG GAA CAA ATG GAG TGT AGG TGT 576
Leu Val Ser Ile Pro Val Asn Pro Lys Glu Gln Met Glu Cys Arg Cys
      180      185      190
GAA AAT GCT GAT GAA GAG GTT GCA ATG GAA GAG GAA GAA GAA GAG 624
Glu Asn Ala Asp Glu Glu Val Ala Met Glu Glu Glu Glu Glu Glu Glu
      195      200      210
GAG GAG GAG GAG GAA GAG GAA ATG GGA AAC CCG GAT GGC TTC TCA CCT 672
Glu Glu Glu Glu Glu Glu Met Gly Asn Pro Asp Gly Phe Ser Pro
      220      225      230
TAG

```

675

- (2) INFORMATION FOR SEQUENCE ID NO: 3:  
    (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 228 base pairs  
        (B) TYPE: nucleic acid  
        (D) TOPOLOGY: linear  
    (ii) MOLECULE TYPE: genomic DNA  
    (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GCATGCAGTT	GCAAAGCCCA	GAAGAAAGAA	ATGGACAGCG	GAAGAAGTGG	TTGTTTTTTT	60
TTCCCCTTCA	TTAATTTTCT	AGTTTTTAGT	AATCCAGAAA	ATTTGATTTT	GTTCTAAAGT	120
TCATTATGCA	AAGATGTCAC	CAACAGACTT	CTGACTGCAT	GGTGAAC TTT	CATATGATAC	180
ATAGGATTAC	ACTTGTACCT	GTAAAAATA	AAAGTTTGAC	TTGCATAC		228

- (2) INFORMATION FOR SEQUENCE ID NO: 4:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 1365 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

ACCACAGGAG AATGAAAAGA ACCCGGGACT CCCAAAGACG CTAGATGTGT      50
GAAGATCCTG ATCACTCATT GGGTGTCTGA GTTCTGCGAT ATTCATCCCT      100
CAGCCAATGA GCTTACTGTT CTCGTGGGGG GTTTGTGAGC CTTGGGTAGG      150
AAGTTTTGCA AGTTCCGCCT ACAGCTCTAG CTTGTGAATT TGTACCCTTT      200
CACGTAAAAA AGTAGTCCAG AGTTTACTAC ACCCTCCCTC CCCCCTCCCA      250
CCTCGTGCTG TGCTGAGTTT AGAAGCTCTC CTTATAGAAG TCTTCCGTAT      300
AGAACTCTTC CGGAGGAAGG AGGGAGGACC CCCCCCTTT GCTCTCCCAG      350
CATGCATTGT GTCAACGCCA TTGCACTGAG CTGGTCGAAG AAGTAAGCCG      400
CTAGCTTGCG ACTCTACTCT TATCTTAACT TAGCTCGGCT TCCTGCTGGT      450
ACCCTTTGTG CC                                         462
ATG TCT GAT AAC AAG AAA CCA GAC AAA GCC CAC AGT GGC TCA      504
GGT GGT GAC GGT GAT GGG AAT AGG TGC AAT TTA TTG CAC CGG      546
TAC TCC CTG GAA GAA ATT CTG CCT TAT CTA GGG TGG CTG GTC      588
TTC GCT GTT GTC ACA ACA AGT TTT CTG GCG CTC CAG ATG TTC      630
ATA GAC GCC CTT TAT GAG GAG CAG TAT GAA AGG GAT GTG GCC      672
TGG ATA GCC AGG CAA AGC AAG CGC ATG TCC TCT GTC GAT GAG      714
GAT GAA GAC GAT GAG GAT GAT GAG GAT GAC TAC TAC GAC GAC      756
GAG GAC GAC GAC GAC GAT GCC TTC TAT GAT GAT GAG GAT GAT      798
GAG GAA GAA GAA TTG GAG AAC CTG ATG GAT GAT GAA TCA GAA      840
GAT GAG GCC GAA GAA GAG ATG AGC GTG GAA ATG GGT GCC GGA      882
GCT GAG GAA ATG GGT GCT GGC GCT AAC TGT GCC TGT GTT CCT      924
GGC CAT CAT TTA AGG AAG AAT GAA GTG AAG TGT AGG ATG ATT      966
TAT TTC TTC CAC GAC CCT AAT TTC CTG GTG TCT ATA CCA GTG     1008
AAC CCT AAG GAA CAA ATG GAG TGT AGG TGT GAA AAT GCT GAT     1050
GAA GAG GTT GCA ATG GAA GAG GAA GAA GAA GAG GAG GAG     1092
GAG GAG GAA GAG GAA ATG GGA AAC CCG GAT GGC TTC TCA CCT     1134
TAG                                                         1137
GCATGCAGTT GCAAAGCCCA GAAGAAAGAA ATGGACAGCG GAAGAAGTGG     1187
TTGTTTTTTT TTCCCCCTCA TTAATTTTCT AGTTTTTGTG AATCCAGAAA     1237
ATTTGATTTT GTTCTAAAGT TCATTATGCA AAGATGTCAC CAACAGACTT     1287
CTGACTGCAT GGTGAACTTT CATATGATAC ATAGGATTAC ACTTGTTACCT     1337
GTTAAAAATA AAAGTTTGAC TTGCATAC                               1365

```



- (2) INFORMATION FOR SEQUENCE ID NO: 5:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 4698 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

```

ACCACAGGAG AATGAAAAGA ACCCGGGACT CCCAAAGACG CTAGATGTGT      50
GAAGATCCTG ATCACTCATT GGGTGTCTGA GTTCTGCGAT ATTCATCCCT      100
CAGCCAATGA GCTTACTGTT CTCGTGGGGG GTTTGTGAGC CTTGGGTAGG      150
AAGTTTTTGA AGTTCCGCCCT ACAGCTCTAG CTTGTGAATT TGTACCCTTT      200
CACGTAAGAAA AGTAGTCCAG AGTTTACTAC ACCCTCCCTC CCCCCTCCCA      250
CCTCGTGCTG TGCTGAGTTT AGAAGTCTTC CTTATAGAAG TCTTCCGTAT      300
AGAAGTCTTC CGGAGGAAGG AGGGAGGACC CCCCCCTTTT GCTCTCCAG      350
CATGCATTGT GTCAACGCCA TTGCACTGAG CTGGTCGAAG AAGTAAGCCG      400
CTAGCTTGCG ACTCTACTCT TATCTTAACT TAGCTCGGCT TCCTGCTGGT      450
ACCCTTTGTG CC                                     462
ATG TCT GAT AAC AAG AAA CCA GAC AAA GCC CAC AGT GGC TCA      504
GGT GGT GAC GGT GAT GGG AAT AGG TGC AAT TTA TTG CAC CGG      546
TAC TCC CTG GAA GAA ATT CTG CCT TAT CTA GGG TGG CTG GTC      588
TTC GCT GTT GTC ACA ACA AGT TTT CTG GCG CTC CAG ATG TTC      630
ATA GAC GCC CTT TAT GAG GAG CAG TAT GAA AGG GAT GTG GCC      672
TGG ATA GCC AGG CAA AGC AAG CGC ATG TCC TCT GTC GAT GAG      714
GAT GAA GAC GAT GAG GAT GAT GAG GAT GAC TAC TAC GAC GAC      756
GAG GAC GAC GAC GAC GAT GCC TTC TAT GAT GAT GAG GAT GAT      798
GAG GAA GAA GAA TTG GAG AAC CTG ATG GAT GAT GAA TCA GAA      840
GAT GAG GCC GAA GAA GAG ATG AGC GTG GAA ATG GGT GCC GGA      882
GCT GAG GAA ATG GGT GCT GGC GCT AAC TGT GCC T          916
GTGAGTAACC CGTGGTCTTT ACTCTAGATT CAGGTGGGGT GCATTCTTTA      966
CTCTTGCCCA CATCTGTAGT AAAGACCACA TTTTGGTTGG GGGTCATTGC     1016
TGGAGCCATC CTTGGCTCTC CTGTCCACGC CTATCCCCGC TCCTCCCATC     1066
CCCCACTCCT TGCTCCGCTC TCTTTCCTTT TCCCACCTTG CCTCTGGAGC     1116
TTCAGTCCAT CTTGCTCTGC TCCCTTTCCC CTTTGTCTCTC CTTGCTCCCC     1166
TCCCCCTCGG CTCAACTTTT CGTGCCTTCT GCTCTCTGAT CCCCACCTC      1216
TTCAGGCTTC CCCATTGCTT CCTCTCCCGA AACCTCCCTC TTCCTGTTCC      1266
CCTTTTCGCG CCTTTTCTTT CCTGCTCCCC TCCCCCTCCC TATTTACCTT      1316
TCACCACTTT TGCTCTCCCT GCTCCCCCTC CCCTTTTGCA CCTTTTCTTT      1366
TCCTGCTCCC CTCCCCCTCC CCTCCCTGTT TACCCTTAC CGCTTTTCCT      1416
CTACCTGCTT CCCTCCCCCT TGCTGCTCCC TCCCTATTTG CATTTTCGGG      1466
TGCTCCTCCC TCCCCCTCCC CCTCCCTCCC TATTTGCATT TTCGGGTGCT      1516
CCTCCCCTCC CCTCCCCAGG CCTTTTTTTT TTTTTTTTTT TTTTTTTTTT      1566
TTGGTTTTTC GAGACAGGTT TTCTCTTTGT ATCCCTGGCT GTCCTGGCAC      1616
TCACTCTGTA GACCAGGCTG GCCTCAAAC T CAGAAATCTG CCTGCCTCTG      1666
CCTCCCAAAT GCTGGGATTA AAGGCTTGCA CCAGGACTGC CCCAGTGCAG      1716
GCCTTTCTTT TTTCTCCTCT CTGGTCTCCC TAATCCCTTT TCTGCATGTT      1766
AACTCCCCCT TTGGCACCTT TCCTTTACAG GACCCCTCC CCTCCCTGT      1816
TTCCCTTCCG GCACCTTCC TAGCCCTGCT CTGTTCCCTC TCCCTGCTCC      1866
CCTCCCCCTC TTTGCTCGAC TTTTAGCAGC CTTACCTCTC CCTGCTTTCT      1916
GCCCGGTTCC CCTTTTTTGT GCCTTTCCTC CTGGCTCCCC TCCACCTTCC      1966
AGCTCACCTT TTTGTTTGTG TGTTTGTGTT GTTGTGTTGTT TGCTTTTTTT      2016
TTTTTTTTTT GCACCTTGTT TTCCAAGATC CCCCTCCCCC TCCGGCTTCC      2066
CCTCTGTGTG CCTTTCTCTG TCCCTCCCCC TCGCTGGCTC CCCCTCCCTT      2116
TCTGCTTTTC CTGTCCCTGC TCCCTTCTCT GCTAACCTTT TAATGCCTTT      2166
CTTTTCTAGA CTCCCCCTC CAGGCTTGCT GTTTGCTTCT GTGCACTTTT      2216
CCTGACCTG CTCCCCCTCC CCTCCCAGCT CCCCCCTCTT TCCCCACCTC      2266
CCTTTCTCCA GCCTGTCACC CCTCCTTCTC TCCTCTCTGT TTCTCCCACT      2316
TCCTGCTTCC TTTACCCCTT CCTCTCCCT ACTCTCCTCC CTGCTGCTG      2366
GACTTCTCT CCAGCCGCCC AGTTCCCTGC AGTCCCTGGAG TCTTTCCTGC      2416
CTCTCTGTCC ATCACTTCCC CCTAGTTTCA CTTCCCTTTT ACTCTCCCT      2466
ATGTGTCTCT CTTCTATCT ATCCCTTCTT TTCTGTCCCC TCTCCTCTGT      2516
CCATCACCTC TCTCCTCCCT TCCCTTTCCT CTCTCTTCCA TTTTCTTCCA      2566
CCTGCTTCTT TACCCTGCCT CTCCCATTGC CCTCTTACCT TTATGCCCCAT      2616
TCCATGTCCC CTCTCAATTC CCGTGTCCCAT TGTGCTCCCT CACATCTTCC      2666

```

ATTTCCCTCT	TTCTCCCTTA	GCCTCTTCTT	CCTCTTCTCT	TGTATCTCCC	2716
TTCCCTTTGC	TTCTCCCTCC	TCCTTTCCCC	TTCCCTATG	CCCTCTACTC	2766
TACTTGATCT	TCTCTCCTCT	CCACATACCC	TTTTTCCTTT	CCACCCTGCC	2816
CTTTGTCCCC	AGACCCTACA	GTATCCTGTG	CACAGGAAGT	GGGAGGTGCC	2866
ATCAACAACA	AGGAGGCAAG	AAACAGAGCA	AAATCCCCAA	ATCAGCAGGA	2916
AAGGCTGGAT	GAAAATAAGG	CCAGGTTCTG	AGGACAGCTG	GAATCTAGCC	2966
AAGTGGCTCC	TATAACCCTA	AGTACCAAGG	GAGAAAAGTGA	TGGTGAAGTT	3016
CTTGATCCTT	GCTGCTTCTT	TTACATATGT	TGGCACATCT	TTCTCAAATG	3066
CAGGCCATGC	TCCATGCTTG	GCGCTTGCTC	AGCGTGGTTA	AGTAATGGGA	3116
GAATCTGAAA	ACTAGGGGCC	AGTGGTTTGT	TTGGGGGACA	AATTAGCACG	3166
TAGTGATATT	TCCCCCTAAA	AATTATAACA	AACAGATTCA	TGATTTGAGA	3216
TCCTTCTACA	GGTGAGAAGT	GGAAAAATTG	TCACTATGAA	GTTCTTTTGA	3266
GGCTAAAGAT	ACTTGGAACC	ATAGAAGCGT	TGTTAAAATA	CTGCTTTCTT	3316
TTGCTAAAA	ATTCTTTCTC	ACATATTCAT	ATTCTCCAG		3355
GT GTT CCT	GGC CAT CAT	TTA AGG	AAG AAT GAA	GTG AAG TGT	3396
AGG ATG ATT	TAT TTC TTC	CAC GAC	CCT AAT TTC	CTG GTG TCT	3438
ATA CCA GTG	AAC CCT AAG	GAA CAA	ATG GAG TGT	AGG TGT GAA	3480
AAT GCT GAT	GAA GAG GTT	GCA ATG	GAA GAG GAA	GAA GAA GAA	3522
GAG GAG GAG	GAG GAG GAA	ATG GGA	AAC CCG	GAT GGC	3564
TTC TCA CCT	TAG				3576
GCATGCAGGT	ACTGGCTTCA	CTAACCAACC	ATTCCTAACA	TATGCCTGTA	3626
GCTAAGAGCA	TCTTTTAAAA	AAATATTATT	GGTAAACTAA	ACAATTGTGA	3676
TCTTTTACAC	TTAATAAGTA	TTAAATTAAT	CCAGTATACA	GTTTAAAGAA	3726
CCCTAAGTTA	AACAGAAGTC	AATGATGTCT	AGATGCCTGT	TCTTTAGATT	3776
GTAGTGAGAC	TACTTACTAC	AGATGAGAAG	TTGTTAGACT	CGGGAGTAGA	3826
GACCAGTAAA	AGATCATGCA	GTGAAATGTG	GCCATGGAAA	TCGCATATTG	3876
TTCTTATAGT	ACCTTTGAGA	CAGCTGATAA	CAGCTGACAA	AAATAAGTGT	3926
TTCAAGAAAG	ATCACACGCC	ATGGTTCACA	TGCAAATTAT	TATTTTGTCT	3976
TTCTGATTTT	TTTCATTTCT	AGACCTGTGG	TTTTAAAGAG	ATGAAAATCT	4026
CTTAAATTTT	CCTTCATCTT	TAATTTTCCT	TAACCTTAGT	TTTTTTCCT	4076
TAGAATTCAA	TTCAAATTCT	TAATTCAATC	TTAATTTTGA	GATTTCTTAA	4126
AATGTTTTTT	AAAAAAAATG	CAAATCTCAT	TTTTAAGAGA	TGAAAGCAGA	4176
GTAAGTGGGG	GGCTTAGGGA	ATCTGTAGGG	TTGCGGTATA	GCAATAGGGA	4226
GTTCTGGTCT	CTGAGAAGCA	GTGAGAGAGA	ATGGAAAACC	AGGCCCTTGC	4276
CAGTAGGTTA	GTGAGGTTGA	TATGATCAGA	TTATGGACAC	TCTCCAAATC	4326
ATAAATACTC	TAACAGCTAA	GGATCTCTGA	GGGAAACACA	ACAGGGAAAT	4376
ATTTTAGTTT	CTCCTTGAGA	AACAATGACA	AGACATAAAA	TTGGCAAGAA	4426
AGTCAGGAGT	GTATTCTAAT	AAGTGTGCT	TATCTCTTAT	TTTCTTCTAC	4476
AGTTGCAAG	CCCAGAAGAA	AGAAATGGAC	AGCGGAAGAA	GTGGTTGTTT	4526
TTTTTTCCCC	TTCAATTAAT	TTCTAGTTTT	TAGTAATCCA	GAAAATTTGA	4576
TTTTGTCTTA	AAGTTCATTA	TGCAAGATG	TCACCAACAG	ACTTCTGACT	4626
GCATGGTGAA	CTTTCATATG	ATACATAGGA	TTACACTTGT	ACCTGTAAAA	4676
AATAAAAGTT	TGACTTGCAAT	AC			4698

- (2) INFORMATION FOR SEQUENCE ID NO: 6:  
(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 9 amino acids  
    (B) TYPE: amino acid  
    (D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: protein  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Leu Pro Tyr Leu Gly Trp Leu Val Phe  
5

- (2) INFORMATION FOR SEQUENCE ID NO: 7:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 2418 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GGATCCAGGC	CCTGCCAGGA	AAAATATAAG	GGCCCTGCGT	GAGAACAGAG	50
GGGGTCATCC	ACTGCATGAG	AGTGGGGATG	TCACAGAGTC	CAGCCCACCC	100
TCCTGGTAGC	ACTGAGAAGC	CAGGGCTGTG	CTTGCGGTCT	GCACCCTGAG	150
GGCCCGTGGA	TTCCTCTTCC	TGGAGCTCCA	GGAAACCAGG	AGTGAGGCCT	200
TGGTCTGAGA	CAGTATCCTC	AGGTCACAGA	GCAGAGGATG	CACAGGGTGT	250
GCCAGCAGTG	AATGTTTGCC	CTGAATGCAC	ACCAAGGGCC	CCACCTGCCA	300
CAGGACACAT	AGGACTCCAC	AGAGTCTGGC	CTCACCTCCC	TACTGTCACT	350
CCTGTAGAAT	CGACCTCTGC	TGGCCGGCTG	TACCCTGAGT	ACCCTCTCAC	400
TTCTCTCCTT	AGGTTTTTCAG	GGGACAGGCC	AACCCAGAGG	ACAGGATTCC	450
CTGGAGGCCA	CAGAGGAGCA	CCAAGGAGAA	GATCTGTAAG	TAGGCCTTTG	500
TTAGAGTCTC	CAAGGTTTCAG	TTCTCAGCTG	AGGCCTCTCA	CACACTCCCT	550
CTCTCCCCAG	GCCTGTGGGT	CTTCATTGCC	CAGCTCCTGC	CCACACTCCT	600
GCCTGCTGCC	CTGACGAGAG	TCATCATGTC	TCTTGAGCAG	AGGAGTCTGC	650
ACTGCAAGCC	TGAGGAAGCC	CTTGAGGCC	AACAAGAGGC	CCTGGGCCTG	700
GTGTGTGTGC	AGGCTGCCAC	CTCCTCCTCC	TCTCCTCTGG	TCCTGGGCAC	750
CCTGGAGGAG	GTGCCCCTG	CTGGGTCAAC	AGATCCTCCC	CAGAGTCCCT	800
AGGGAGCCTC	CGCCTTTCCC	ACTACCATCA	ACTTCACTCG	ACAGAGGCAA	850
CCCAGTGAGG	GTTCCAGCAG	CCGTGAAGAG	GAGGGGCCAA	GCACCTCTTG	900
TATCCTGGAG	TCCTTGTTCC	GAGCAGTAAT	CACCTAAGAAG	GTGGCTGATT	950
TGGTTGGTTT	TCTGCTCCTC	AAATATCGAG	CCAGGGAGCC	AGTCACAAAG	1000
GCAGAAATGC	TGGAGAGTGT	CATCAAAAAT	TACAAGCACT	GTTTTCTCTGA	1050
GATCTTCCGC	AAAGCCTCTG	AGTCCTTGCA	GCTGGTCTTT	GGCATTGACG	1100
TGAAGGAAGC	AGACCCACCC	GGCCACTCCT	ATGTCCTTGT	CACCTGCCTA	1150
GGTCTCTCCT	ATGATGGCCT	GCTGGGTGAT	AATCAGATCA	TGCCCCAGAC	1200
AGGCTTCCTG	ATAATTGTCC	TGGTCATGAT	TGCAATGGAG	GGCGGCCATG	1250
CTCCTGAGGA	GGAAATCTGG	GAGGAGCTGA	GTGTGATGGA	GGTGTATGAT	1300
GGGAGGGAGC	ACAGTGCCCTA	TGGGGAGCCC	AGGAAGCTGC	TCACCCAAGA	1350
TTTGGTGCGA	GAAAAGTACC	TGGAGTACGG	CAGGTGCCGG	ACAGTGATCC	1400
CGCAGCGTAT	GAGTTCCTGT	GGGGTCCAAG	GGCCCTCGCT	GAAACCAGCT	1450
ATGTGAAAGT	CCTTGAGTAT	GTGATCAAGG	TCAGTGCAAG	AGTTCGCTTT	1500
TTCTTCCCAT	CCCTGCGTGA	AGCAGCTTTG	AGAGAGGAGG	AAGAGGGAGT	1550
CTGAGCATGA	GTTGCAGCCA	AGGCCAGTGG	GAGGGGGACT	GGGCCAGTGC	1600
ACCTTCCAGG	GCCGCGTCCA	GCAGCTTCCC	CTGCCTCGTG	TGACATGAGG	1650
CCCATTCTTC	ACTCTGAAGA	GAGCGGTCAG	TGTTCTCAGT	AGTAGGTTTC	1700
TGTTCTATTG	GGTGACTTGG	AGATTTATCT	TTGTTCTCTT	TTGGAATTGT	1750
TCAAATGTTT	TTTTTTAAGG	GATGGTTGAA	TGAACTTCAG	CATCCAAGTT	1800
TATGAATGAC	AGCAGTCACA	CAGTTCTGTG	TATATAGTTT	AAGGGTAAGA	1850
GTCTTGTGTT	TTATTTCAGAT	TGGGAAATCC	ATTCTATTTT	GTGAATTGGG	1900
ATAATAACAG	CAGTGGAATA	AGTACTTAGA	AATGTGAAAA	ATGAGCAGTA	1950
AAATAGATGA	GATAAAGAAC	TAAAGAAAAT	AAGAGATAGT	CAATTCCTGC	2000
CTTATACCTC	AGTCTATTCT	GTAAAAATTT	TAAAGATATA	TGCATACCTG	2050
GATTTCTCTG	GCTTCTTTGA	GAATGTAAAG	GAAATTAAT	CTGAATAAAG	2100
AATTCCTCCT	GTTCACTGGC	TCTTTTCTTC	TCCATGCACT	GAGCATCTGC	2150
TTTTTGGGAAG	GCCCTGGGTT	AGTAGTGGAG	ATGCTAAGGT	AAGCCAGACT	2200
CATACCCACC	CATAGGGTCG	TAGAGTCTAG	GAGCTGCAGT	CACGTAATCG	2250
AGGTGGCAAG	ATGTCCTCTA	AAGATGTAGG	GAAAAGTGAG	AGAGGGGTGA	2300
GGGTGTGGGG	CTCCGGGTGA	GAGTGGTGGA	GTGTCAATGC	CCTGAGCTGG	2350
GGCATTTTGG	GCTTTGGGAA	ACTGCAGTTC	CTTCTGGGGG	AGCTGATTGT	2400
AATGATCTTG	GGTGGATCC				2418

- (2) INFORMATION FOR SEQUENCE ID NO: 8:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 5724 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: MAGE-1 gene  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

CCCCGGGGCAC	CACTGGCATC	CCTCCCCCTA	CCACCCCA	TCCCTCCCTT	50
TACGCCACCC	ATCCAAACAT	CTTCACGCTC	ACCCCCAGCC	CAAGCCAGGC	100
AGAATCCGGT	TCCACCCCTG	CTCTCAACCC	AGGGAAGCCC	AGGTGCCCAG	150
ATGTGACGCC	ACTGACTTGA	GCATTAGTGG	TTAGAGAGAA	GCGAGGTTTT	200
CGGTCTGAGG	GGCGGCTTGA	GATCGGTGGA	GGGAAGCGGG	CCAGCTCTG	250
TAAGGAGGCA	AGGTGACATG	CTGAGGGAGG	ACTGAGGACC	CACTTACCCC	300
AGATAGAGGA	CCCCAAATAA	TCCCTTCATG	CCAGTCCTGG	ACCATCTGGT	350
GGTGGACTTC	TCAGGCTGGG	CCACCCCCAG	CCCCCTTGCT	GCTTAAACCA	400
CTGGGGACTC	GAAGTCAGAG	CTCCGTGTGA	TCAGGGAAGG	GCTGCTTAGG	450
AGAGGGCAGC	GTCCAGGCTC	TGCCAGACAT	CATGCTCAGG	ATTCTCAAGG	500
AGGGCTGAGG	GTCCCTAAGA	CCCCACTCCC	GTGACCCAAC	CCCCACTCCA	550
ATGCTCACTC	CCGTGACCCA	ACCCCTCTTT	CATTGTCTAT	CCAACCCCCA	600
CCCCACATCC	CCCACCCCAT	CCCTCAACCC	TGATGCCCAT	CGCCCCAGCC	650
ATTCCACCTT	CACCCCCACC	CCCACCCCCA	CGCCCACTCC	CACCCCCACC	700
CAGGCAGGAT	CCGGTTCCCG	CCAGGAAACA	TCCGGGTGCC	CGGATGTGAC	750
GCCACTGACT	TGCGCATTGT	GGGGCAGAGA	GAAGCGAGGT	TTCCATTCTG	800
AGGGACGCGG	TAGAGTTCGG	CCGAAGGAAC	CTGATGCCAT	CTCTGTGAGG	850
AGGCAAGGTG	AGAGGCTGAG	GGAGGACTGA	GGACCCCGCC	ACTCCAAATA	900
GAGAGCCCCA	AATATTCCAG	CCCCGCCCTT	GCTGCCAGCC	CTGGCCCAAC	950
CGCGGGAAGA	CGTCTCAGCC	TGGGCTGCCC	CCAGACCCCT	GCTCCAAAAG	1000
CCTTGAGAGA	CACCAGGTTT	TTCTCCCCAA	TCTGTGGAAT	CAGAGGTTGC	1050
TGTGACCAGG	GCAGGACTGG	TTAGGAGAGG	GCAGGGCACA	GGCTCTGCCA	1100
GGCATCAAGA	TCAGCACCCA	AGAGGGAGGG	CTGTGGGCCC	CCAAGACTGC	1150
ACTCCAATCC	CCACTCCAC	CCCATTGCGA	TTCCCATTTT	CCACCCAACC	1200
CCCATCTCCT	CAGCTACACC	TCCACCCCCA	TCCCTACTCC	TACTCCGTCA	1250
CCTGACCACC	ACCCTCCAGC	CCCAGCACCA	GCCCCAACCC	TTCTGCCACC	1300
TCACCTCAC	TGCCCCCAAC	CCCACCTTCA	TCTCTCTCAT	GTGCCCCACT	1350
CCCATCGCCT	CCCCCATTTT	GGCAGAATCC	GGTTTGCCCC	TGCTCTCAAC	1400
CCAGGGAAGC	CCTGGTAGGC	CCGATGTGAA	ACCCTGACT	TGAACCTCAC	1450
AGATCTGAGA	GAAGCCAGGT	TCATTTAATG	GTTCTGAGGG	GCGGCTTGAG	1500
ATCCACTGAG	GGGAGTGGTT	TTAGGCTCTG	TGAGGAGGCA	AGGTGAGATG	1550
CTGAGGGAGG	ACTGAGGAGG	CACACACCCC	AGGTAGATGG	CCCCAAAATG	1600
ATCCAGTACC	ACCCCTGCTG	CCAGCCCTGG	ACCACCCGGC	CAGGACAGAT	1650
GTCTCAGCTG	GACCACCCCC	CGTCCCGTCC	CACTGCCACT	TAACCCACAG	1700
GGCAATCTGT	AGTCATAGCT	TATGTGACCG	GGGCAGGGTT	GGTCAGGAGA	1750
GGCAGGGCCC	AGGCATCAAG	GTCCAGCATC	CGCCCGGCAT	TAGGGTCAGG	1800
ACCCTGGGAG	GGAAGTGAGG	GTTCCCCACC	CACACCTGTC	TCCTCATCTC	1850
CACCGCCACC	CCACTCACAT	TCCCATACCT	ACCCCTTACC	CCCAACCTCA	1900
TCTTGTCAGA	ATCCCTGCTG	TCAACCCACG	GAAGCCACGG	GAATGGCGGC	1950
CAGGCACTCG	GATCTTGACG	TCCCATCCA	GGGTCTGATG	GAGGGAAGGG	2000
GCTTGAACAG	GGCCTCAGGG	GAGCAGAGGG	AGGGCCCTAC	TGCGAGATGA	2050
GGGAGGCCTC	AGAGGACCCA	GCACCCTAGG	ACACCGCACC	CCTGTCTGAG	2100
ACTGAGGCTG	CCACTTCTGG	CCTCAAGAAT	CAGAACGATG	GGGACTCAGA	2150
TTGCATGGGG	GTGGGACCCA	GGCCTGCAAG	GCTTACGCGG	AGGAAGAGGA	2200
GGGAGGACTC	AGGGGACCTT	GGAATCCAGA	TCAGTGTGGA	CCTCGGCCCT	2250
GAGAGGTCCA	GGGCACGGTG	GCCACATATG	GCCCATATTT	CCTGCATCTT	2300
TGAGGTGACA	GGACAGAGCT	GTGGTCTGAG	AAGTGGGGCC	TCAGGTCAAC	2350
AGAGGAGGA	GTTCCAGGAT	CCATATGGCC	CAAGATGTGC	CCCTTCATG	2400
AGGACTGGGG	ATATCCCCGG	CTCAGAAAGA	AGGGACTCCA	CACAGTCTGG	2450
CTGTCCCTTT	TTAGTAGCTC	TAGGGGGACC	AGATCAGGGA	TGGCGGTATG	2500
TTCCATTCTC	ACTTGTACCA	CAGGCAGGAA	GTTGGGGGGC	CCTCAGGGAG	2550
ATGGGCTCTT	GGGGTAAAGG	GGGGATGTCT	ACTCATGTCA	GGGAATTGGG	2600
GGTTGAGGAA	GCACAGGCGC	TGGCAGGAAT	AAAGATGAGT	GAGACAGACA	2650
AGGCTATTGG	AATCCACACC	CCAGAACCAA	AGGGGTGAGC	CCTGGACACC	2700

TCACCCAGGA	TGTGGCTTCT	TTTTCACTCC	TGTTTCCAGA	TCTGGGGCAG	2750
GTGAGGACCT	CATTCTCAGA	GGGTGACTCA	GGTCAACGTA	GGGACCCCCA	2800
TCTGGTCTAA	AGACAGAGCG	GTCCCAGGAT	CTGCCATGCG	TTCGGGTGAG	2850
GAACATGAGG	GAGGACTGAG	GGTACCCAG	GACCAGAACA	CTGAGGGAGA	2900
CTGCACAGAA	ATCAGCCCTG	CCCCTGCTGT	CACCCAGAG	AGCATGGGCT	2950
GGGCCGTCTG	CCGAGGTCCT	TCCGTTATCC	TGGGATCATT	GATGTCAGGG	3000
ACGGGGAGGC	CTTGGTCTGA	GAAGGCTGCG	CTCAGGTCAG	TAGAGGGAGC	3050
GTCCCAGGCC	CTGCCAGGAG	TCAAGGTGAG	GACCAAGCGG	GCACCTCACC	3150
CAGGACACAT	TAATTCGAAT	GAATTTTGAT	ATCTCTTGCT	GCCCTTCCCC	3200
AAGGACCTAG	GCACGTGTGG	CCAGATGTTT	GTCCCCCTCT	GTCCCTCCAT	3250
TCCTTATCAT	GGATGTGAAC	TCTTGATTG	GATTTCTCAG	ACCAGCAAAA	3300
GGGCAGGATC	CAGGCCCTGC	CAGGAAAAAT	ATAAGGGCCC	TGCGTGAGAA	3350
CAGAGGGGGT	CATCCACTGC	ATGAGAGTGG	GGATGTCACA	GAGTCCAGCC	3400
CACCCCTCTG	GTAGCACTGA	GAAGCCAGGG	CTGTGCTTGC	GGTCTGCACC	3450
CTGAGGGCCC	GTGGATTCTT	CTTCCTGGAG	CTCCAGGAAC	CAGGCAGTGA	3500
GGCCTTGCTC	TGAGACAGTA	TCCTCAGGTC	ACAGAGCAGA	GGATGCACAG	3550
GGTGTGCCAG	CAGTGAATGT	TTGCCCTGAA	TGCACACCAA	GGGCCCCACC	3600
TGCCACAGTA	CACATAGGAC	TCCACAGAGT	CTGGCCTCAC	CTCCCTACTG	3650
TCAGTCTGTG	AGAATCGACC	TCTGCTGGCC	GGTGTACCC	TGAGTACCCT	3700
CTCACTTCCT	CCTTCAGGTT	TTCAGGGGAC	AGGCCAACCC	AGAGGACAGG	3750
ATTCCCTGGA	GGCCACAGAG	GAGCACCAAG	GAGAAGATCT	GTAAGTAGGC	3800
CTTTGTTAGA	GTCTCCAAGG	TTCAGTTCTC	AGCTGAGGCC	TCTCACACAC	3850
TCCCTCTCTC	CCCAGGCTTG	TGGGTCTTCA	TTGCCAGCT	CCTGCCCCACA	3900
CTCCTGCCTG	CTGCCCTGAC	GAGAGTCATC			3930
ATG TCT CTT	GAG CAG AGG	AGT CTG CAC	TGC AAG CCT	GAG GAA	3972
GCC CTT GAG	GCC CAA CAA	GAG GCC CTG	GGC CTG GTG	TGT GTG	4014
CAG GCT GCC	ACC TCC TCC	TCT CCT CTG	GTC CTG GGC	ACC	4056
CTG GAG GAG	GTG CCC ACT	GCT GGG TCA	ACA GAT CCT	CCC CAG	4098
AGT CCT CAG	GGA GCC TCC	GCC TTT CCC	ACT ACC ATC	AAC TTC	4140
ACT CGA CAG	AGG CAA CCC	AGT GAG GGT	TCC AGC AGC	CGT GAA	4182
GAG GAG GGG	CCA AGC ACC	TCT TGT ATC	CTG GAG TCC	TTG TTC	4224
CGA GCA GTA	ATC ACT AAG	AAG GTG GCT	GAT TTG GTT	GGT TTT	4266
CTG CTC CTC	AAA TAT CGA	GCC AGG GAG	CCA GTC ACA	AAG GCA	4308
GAA ATG CTG	GAG AGT GTC	ATC AAA AAT	TAC AAG CAC	TGT TTT	4350
CCT GAG ATC	TTC GGC AAA	GCC TCT GAG	TCC TTG CAG	CTG GTC	4392
TTT GGC ATT	GAC GTG AAG	GAA GCA GAC	CCC ACC GGC	CAC TCC	4434
TAT GTC CTT	GTC ACC TGC	CTA GGT CTC	TCC TAT GAT	GGC CTG	4476
CTG GGT GAT	AAT CAG ATC	ATG CCC AAG	ACA GGC TTC	CTG ATA	4518
ATT GTC CTG	GTC ATG ATT	GCA ATG GAG	GGC GGC CAT	GCT CCT	4560
GAG GAG GAA	ATC TGG GAG	GAG CTG AGT	GTG ATG GAG	GTG TAT	4602
GAT GGG AGG	GAG CAC AGT	GCC TAT GGG	GAG CCC AGG	AAG CTG	4644
CTC ACC CAA	GAT TTG GTG	CAG GAA AAG	TAC CTG GAG	TAC GGC	4686
AGG TGC CGG	ACA GTG ATC	CCG CAC GCT	ATG AGT TCC	TGT GGG	4728
GTC CAA GGG	CCC TCG CTG	AAA CCA GCT	ATG TGA		4761
AAGTCCTTGA	GTATGTGATC	AAGGTCAGTG	CAAGAGTTC		4800
GCTTTTTCTT	CCCATCCCTG	CGTGAAGCAG	CTTTGAGAGA	GGAGGAAGAG	4850
GGAGTCTGAG	CATGAGTTGC	AGCCAAGGCC	AGTGGGAGGG	GGACTGGGCC	4900
AGTGCACCTT	CCAGGGCCGC	GTCCAGCAGC	TTCCCCTGCC	TGCTGTGACA	4950
TGAGGCCCCAT	TCTTCACTCT	GAAGAGAGCG	GTCAGTGTTT	TCAGTAGTAG	5000
GTTTCTGTTC	TATTGGGTGA	CTTGGAGATT	TATCTTTGTT	CTCTTTTGGA	5050
ATTGTTCAAA	TGTTTTTTTT	TAAGGGATGG	TTGAATGAAC	TTCAGCATCC	5100
AAGTTTATGA	ATGACAGCAG	TCACACAGTT	CTGTGTATAT	AGTTTAAGGG	5150
TAAGAGTCTT	GTGTTTTTAT	CAGATTGGGA	AATCCATTCT	ATTTTGTGAA	5200
TTGGGATAAT	AACAGCAGTG	GAATAAGTAC	TTAGAAATGT	GAAAAATGAG	5250
CAGTAAAAATA	GATGAGATAA	AGAATAAGAG	AAATTAAGAG	ATAGTCAATT	5300
CTTGCCCTTAT	ACCTCAGTCT	ATTCTGTAAA	ATTTTAAAG	ATATATGCAT	5350
ACCTGGATTT	CCTTGGCTTC	TTTGAGAAATG	TAAGAGAAAT	TAAATCTGAA	5400
TAAAGAATTG	TTCTGTCTCA	CTGGCTCTTT	TCTTCTCCAT	GCACTGAGCA	5450
TCTGCTTTTT	GGAAGGCCCT	GGGTAGTAG	TGGAGATGCT	AAGGTAAGCC	5500
AGACTCATAC	CCACCCATAG	GGTCGTAGAG	TCTAGGAGCT	GCAGTCACGT	5550
AATCGAGGTG	GCAAGATGTC	CTCTAAAGAT	GTAGGGAAAA	GTGAGAGAGG	5600
GGTGAGGGTG	TGGGGCTCCG	GGTGAGAGTG	GTTGAGTGTG	AATGCCCTGA	5650
GCTGGGGCAT	TTTGGGCTTT	GGGAAACTGC	AGTTCTTCTT	GGGGGAGCTG	5700
ATTGTAATGA	TCTTGGGTGG	ATCC			5724

- (2) INFORMATION FOR SEQUENCE ID NO: 9:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 4157 base pairs
    - (B) TYPE: nucleic acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: genomic DNA
  - (ix) FEATURE:
    - (A) NAME/KEY: MAGE-2 gene
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

```

CCCATCCAGA TCCCCATCCG GGCAGAATCC GGTTCACCC TTGCCGTGAA      50
CCGAGGGAAG TCACGGGCCC GGATGTGACG CCACTGACTT GCACATTGGA      100
GGTCAGAGGA CAGCGAGATT CTCGCCCTGA GCAACGGCCT GACGTCGGCG      150
GAGGGAAGCA GGCGCAGGCT CCGTGAGGAG GCAAGGTAAG ACGCCGAGGG      200
AGGACTGAGG CGGGCCTCAC CCCAGACAGA GGGCCCCCAA TTAATCCAGC      250
GCTGCCCTCTG CTGCCGGGCC TGGACCACCC TGCAGGGGAA GACTTCTCAG      300
GCTCAGTCGC CACCACCTCA CCCCGCCACC CCCCGCCGCT TTAACCGCAG      350
GGAACCTCTGG CGTAAGAGCT TTGTGTGACC AGGGCAGGGC TGGTTAGAAG      400
TGCTCAGGGC CCAGACTCAG CCAGGAATCA AGGTCAGGAC CCCAAGAGGG      450
GACTGAGGGC AACCACCCCC CTACCCTCAC TACCAATCCC ATCCCCAAC      500
ACCAACCCCC CCCCCATCCC TCAAACACCA ACCCACCCCC CAAACCCCAT      550
TCCCATCTCC TCCCCACCA CCATCCTGGC AGAATCCGGC TTTGCCCTG      600
CAATCAACCC ACGGAAGCTC CGGGAATGGC GGCCAAGCAC GCGGATCCTG      650
ACGTTACATC GTACGGCTAA GGGAGGGAAG GGGTTGGGTC TCGTGAGTAT      700
GGCCTTTGGG ATGCAGAGGA AGGGCCAGG CCACTTGAA GACAGTGGAG      750
TCCTTAGGGG ACCCAGCATG CCAGGACAGG GGGCCCACTG TACCCCTGTC      800
TCAAACCTGAG CCACCTTTTC ATTACAGCCGA GGAATCCTA GGGATGCAGA      850
CCCACTTCAG GGGGTTGGGG CCCAGCCTGC GAGGAGTCAA GGGGAGGAAG      900
AAGAGGGGAG ACTGAGGGGA CCTTGGAGTC ACAGTCAGTG GCAACCTTGG      950
GCTGGGGGAT CCTGGGCACA GTGGCCGAAT GTGCCCCGTG CTCATTGCAC      1000
CTTCAGGGTG ACAGAGAGTT GAGGGCTGTG GTCTGAGGGC TGGGACTTCA      1050
GGTCAGCAGA GGGAGGAATC CCAGGATCTG CCGGACCCAA GGTGTGCCCC      1100
CTTCATGAGG ACTCCCCATA CCCCCGGCCC AGAAGAAGG GATGCCACAG      1150
AGTCTGGAAG TAAATTGTTC TTAGCTCTGG GGAACCTGA TCAGGGATGG      1200
CCCTAAGTGA CAATCTCATT TGTACCACAG GCAGGAGGTT GGGGAACCCCT      1250
CAGGCAGATA AGGTGTTGGT GTAAAGAGGA GCTGTCTGCT CATTTAGGG      1300
GGTTCCCCCT TGAGAAAGGG CAGTCCCTGG CAGGAGTAAA GATGAGTAAC      1350
CCACAGGAGG CCATCATAAC GTTCACCCTA GAACCAAAGG GGTGAGCCCT      1400
GGACAACGCA CGTGGGGTAA CAGGATGTGG CCCCTCCTCA CTTGTCTTTC      1450
CAGATCTCAG GGAGTTGATG ACCTTGTTTT CAGAAGGTGA CTCAGTCAAC      1500
ACAGGGGGCCC CTCTGGTCTGA CAGATGCAGT GGTCTTAGGA TCTGCCAAGC      1550
ATCCAGGTGG AGAGCCTGAG GTAGGATTGA GGGTACCCCT GGGCCAGAAT      1600
GCAGCAAGGG GGCCCCATAG AAATCTGCCC TGCCCCCTCG GTTACTTCAG      1650
AGACCCTGGG CAGGGCTGTC AGCTGAAGTC CCTCCATTAT CTGGGATCTT      1700
TGATGTCAGG GAAGGGGAGG CTTTGGTCTG AAGGGGCTGG AGTCAGGTCA      1750
GTAGAGGGAG GGTCTCAGGC CCTGCCAGGA GTGGACGTGA GGACCAAGCG      1800
GACTCGTCAC CCAGGACACC TGGACTCCAA TGAATTGAC ATCTCTCGTT      1850
GTCTTTCGCG GAGGACCTGG TCACGTATGG CCAGATGTGG GTCCCCCTTA      1900
TCTCCTTCTG TACCATATCA GGGATGTGAG TTCTTGACAT GAGAGATTCT      1950
CAAGCCAGCA AAAGGGTGGG ATTAGGCCCT ACAAGGAGAA AGGTGAGGGC      2000
CCTGAGTGAG CACAGAGGGG ACCCTCCACC CAAGTAGAGT GGGGACCTCA      2050
CGGAGTCTGG CCAACCCTGC TGAGACTTCT GGAATCCGT GGCTGTGCTT      2100
GCAGTCTGCA CACTGAAGGC CCGTGCAATC CTCTCCAGG AATCAGGAGC      2150
TCCAGGAACC AGGCAGTGAG GCCTTGGTCT GAGTCAGTGC CTCAGGTCAC      2200
AGAGCAGAGG GGACGCAGAG AGTGCCAACA CTGAAGGTTT GCCTGGAATG      2250
CACACCAAGG GCCCAACCCG CCCAGAACAA ATGGGACTCC AGAGGGCCTG      2300
GCCTCACCCCT CCCTATTCTC AGTCCTGCAG CCTGAGCATG TGCTGGCCGG      2350
CTGTACCCTG AGGTGCCCTC CCACTTCCTC CTTTCAGTTC TGAGGGGGAC      2400
AGGCTGACAA GTAGGACCCG AGGCACTGGA GGAGCATTGA AGGAGAAGAT      2450
CTGTAAGTAA GCCTTTGTCA GAGCCTCCAA GGTTTCAGTT AGTTCTCACC      2500
TAAGGCCTCA CACACGCTCC TTCTCTCCCC AGGCCTGTGG GTCTTCATTG      2550
CCCAGCTCCT GCGCCGACTC CTGCCTGCTG CCCTGACCAG AGTCATC      2597
ATG CCT CTT GAG CAG AGG AGT CAG CAC TGC AAG CCT GAA GAA      2639
GGC CTT GAG GCC CGA GGA GAG GCC CTG GGC CTG GTG GGT GCG      2681
CAG GCT CCT GCT ACT GAG GAG CAG CAG ACC GCT TCT TCC TCT      2723

```

TCT ACT CTA GTG GAA GTT ACC CTG GGG GAG GTG CCT GCT GCC	2765
GAC TCA CCG AGT CCT CCC CAC AGT CCT CAG GGA GCC TCC AGC	2807
TTC TCG ACT ACC ATC AAC TAC ACT CTT TGG AGA CAA TCC GAT	2849
GAG GGC TCC AGC AAC CAA GAA GAG GAG GGG CCA AGA ATG TTT	2891
CCC GAC CTG GAG TCC GAG TTC CAA GCA GCA ATC AGT AGG AAG	2933
ATG GTT GAG TTG GTT CAT TTT CTG CTC CTC AAG TAT CGA GCC	2975
AGG GAG CCG GTC ACA AAG GCA GAA ATG CTG GAG AGT GTC CTC	3017
AGA AAT TGC CAG GAC TTC TTT CCC GTG ATC TTC AGC AAA GCC	3059
TCC GAG TAC TTG CAG CTG GTC TTT GGC ATC GAG GTG GTG GAA	3101
GTG GTC CCC ATC AGC CAC TTG TAC ATC CTT GTC ACC TGC CTG	3143
GGC CTC TCC TAC GAT GGC CTG CTG GGC GAC AAT CAG GTC ATG	3185
CCC AAG ACA GGC CTC CTG ATA ATC GTC CTG GCC ATA ATC GCA	3227
ATA GAG GGC GAC TGT GCC CCT GAG GAG AAA ATC TGG GAG GAG	3269
CTG AGT ATG TTG GAG GTG TTT GAG GGG AGG GAG GAC AGT GTC	3311
TTC GCA CAT CCC AGG AAG CTG CTC ATG CAA GAT CTG GTG CAG	3353
GAA AAC TAC CTG GAG TAC CGG CAG GTG CCC GGC AGT GAT CCT	3395
GCA TGC TAC GAG TTC CTG TGG GGT CCA AGG GCC CTC ATT GAA	3437
ACC AGC TAT GTG AAA GTC CTG CAC CAT ACA CTA AAG ATC GGT	3479
GGA GAA CCT CAC ATT TCC TAC CCA CCC CTG CAT GAA CGG GCT	3521
TTG AGA GAG GGA GAA GAG TGA	3542
GTCTCAGCAC ATGTTGCAGC CAGGGCCAGT GGGAGGGGGT CTGGGCCAGT	3592
GCACCTTCCA GGGCCCCATC CATTAGCTTC CACTGCCTCG TGTGATATGA	3642
GGCCCATTC TGCCTCTTTG AAGAGAGCAG TCAGCATTCT TAGCAGTGAG	3692
TTTCTGTTCT GTTGGATGAC TTTGAGATTT ATCTTTCTTT CCTGTTGGAA	3742
TTGTTCAAAT GTTCCTTTTA ACAAATGGTT GGATGAACTT CAGCATCCAA	3792
GTTTTATGAAT GACAGTAGTC ACACATAGTG CTGTTTATAT AGTTTAGGGG	3842
TAAGAGTCCT GTTTTTTATT CAGATTGGA AATCCATTCC ATTTTGTGAG	3892
TTGTCACATA ATAACAGCAG TGGAATATGT ATTTGCCTAT ATTGTGAACG	3942
AATTAGCAGT AAAATACATG ATACAAGGAA CTCAAAAGAT AGTTAATTCT	3992
TGCCTTATAC CTCAGTCTAT TATGTAAAAT TAAAAATATG TGTATGTTTT	4042
TGCTTCTTTG AGAATGCAAA AGAAATTAAA TCTGAATAAA TTCTTCCTGT	4092
TCACTGGCTC ATTTCTTTAC CATTCACTCA GCATCTGCTC TGTGGAAGGC	4142
CCTGGTAGTA GTGGG	4157



- (2) INFORMATION FOR SEQUENCE ID NO: 10:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 662 base pairs
    - (B) TYPE: nucleic acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: genomic DNA
  - (ix) FEATURE:
    - (A) NAME/KEY: MAGE-21 gene
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

GGATCCCCAT	GGATCCAGGA	AGAATCCAGT	TCCACCCCTG	CTGTGAACCC	50
AGGGAAGTCA	CGGGGCCGGA	TGTGACGCCA	CTGACTTGCG	CGTTGGAGGT	100
CAGAGAACAG	CGAGATTCTC	GCCCTGAGCA	ACGGCCTGAC	GTCTGGCGGAG	150
GGAAGCAGGC	GCAGGCTCCG	TGAGGAGGCA	AGGTAAGATG	CCGAGGGAGG	200
ACTGAGGCGG	GCCTCACCCC	AGACAGAGGG	CCCCCAATAA	TCCAGCGCTG	250
CCTCTGCTGC	CAGGCCTGGA	CCACCCTGCA	GGGGAAGACT	TCTCAGGCTC	300
AGTCGCCACC	ACCTCACCCC	GCCACCCCCC	GCCGCTTTAA	CCGCAGGGAA	350
CTCTGGTGTA	AGAGCTTTGT	GTGACCAGGG	CAGGGCTGGT	TAGAAGTGCT	400
CAGGGCCCCAG	ACTCAGCCAG	GAATCAAGGT	CAGGACCCCA	AGAGGGGACT	450
GAGGGTAACC	CCCCCGCACC	CCCACCACCA	TTCCCATCCC	CCAACACCAA	500
CCCCACCCCC	ATCCCCCAAC	ACCAAACCCA	CCACCATCGC	TCAAACATCA	550
ACGGCACCCC	CAAACCCCGA	TTCCCATCCC	CACCCATCCT	GGCAGAATCG	600
GAGCTTTGCC	CCTGCAATCA	ACCCACGGAA	GCTCCGGGAA	TGGCGGCCAA	650
GCACGCGGAT	CC				662

- (2) INFORMATION FOR SEQUENCE ID NO: 11:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 1640 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: cDNA to mRNA  
 (ix) FEATURE:  
     (A) NAME/KEY: cDNA MAGE-3  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

```

GCCGCGAGGG AAGCCGGCCC AGGCTCGGTG AGGAGGCAAG GTTCTGAGGG      50
GACAGGCTGA CCTGGAGGAC CAGAGGCCCC CGGAGGAGCA CTGAAGGAGA      100
AGATCTGCCA GTGGGTCTCC ATTGCCCAGC TCCTGCCCAC ACTCCCGCCT      150
GTTGCCCTGA CCAGAGTCAT C                                     171
ATG CCT CTT GAG CAG AGG AGT CAG CAC TGC AAG CCT GAA GAA      213
GGC CTT GAG GCC CGA GGA GAG GCC CTG GGC CTG GTG GGT GCG      255
CAG GCT CCT GCT ACT GAG GAG CAG GAG GCT GCC TCC TCC TCT      297
TCT ACT CTA GTT GAA GTC ACC CTG GGG GAG GTG CCT GCT GCC      339
GAG TCA CCA GAT CCT CCC CAG AGT CCT CAG GGA GCC TCC AGC      381
CTC CCC ACT ACC ATG AAC TAC CCT CTC TGG AGC CAA TCC TAT      423
GAG GAC TCC AGC AAC CAA GAA GAG GAG GGG CCA AGC ACC TTC      465
CCT GAC CTG GAG TCC GAG TTC CAA GCA GCA CTC AGT AGG AAG      507
GTG GCC GAG TTG GTT CAT TTT CTG CTC CTC AAG TAT CGA GCC      549
AGG GAG CCG GTC ACA AAG GCA GAA ATG CTG GGG AGT GTC GTC      591
GGA AAT TGG CAG TAT TTC TTT CCT GTG ATC TTC AGC AAA GCT      633
TCC AGT TCC TTG CAG CTG GTC TTT GGC ATC GAG CTG ATG GAA      675
GTG GAC CCC ATC GGC CAC TTG TAC ATC TTT GCC ACC TGC CTG      717
GGC CTC TCC TAC GAT GGC CTG CTG GGT GAC AAT CAG ATC ATG      759
CCC AAG GCA GGC CTC CTG ATA ATC GTC CTG GCC ATA ATC GCA      801
AGA GAG GGC GAC TGT GCC CCT GAG GAG AAA ATC TGG GAG GAG      843
CTG AGT GTG TTA GAG GTG TTT GAG GGG AGG GAA GAC AGT ATG      885
TTG GGG GAT CCC AAG AAG CTG CTC ACC CAA CAT TTC GTG CAG      927
GAA AAC TAC CTG GAG TAC CGG CAG GTC CCC GGC AGT GAT CCT      969
GCA TGT TAT GAA TTC CTG TGG GGT CCA AGG GCC CTC GTT GAA     1011
ACC AGC TAT GTG AAA GTC CTG CAC CAT ATG GTA AAG ATC AGT     1053
GGA GGA CCT CAC ATT TCC TAC CCA CCC CTG CAT GAG TGG GTT     1095
TTG AGA GAG GGG GAA GAG TGA                                     1116
GTCTGAGCAC GAGTTGCAGC CAGGGCCAGT GGGAGGGGGT CTGGGCCAGT     1166
GCACCTTCCG GGGCCGCATC CCTTAGTTTC CACTGCCTCC TGTGACGTGA     1216
GGCCCATTTCT TCACTCTTTG AAGCGAGCAG TCAGCATTCT TAGTAGTGGG     1266
TTTCTGTTCT GTTGGATGAC TTTGAGATTA TTCTTTGTTT CCTGTTGGAG     1316
TTGTTCAAAT GTTCCTTTTA ACGGATGGTT GAATGAGCGT CAGCATCCAG     1366
GTTTATGAAT GACAGTAGTC ACACATAGTG CTGTTTATAT AGTTTAGGAG     1416
TAAGAGTCTT GttTTTTACT CAAATTgGGA AATCCATTCC ATTTTGTGAA     1466
TTGTGACATA ATAATAGCAG TGGTAAAAGT ATTTGCTTAA AATTGTGAGC     1516
GAATTAGCAA TAACATACAT GAGATAACTC AAGAAATCAA AAGATAGTTG     1566
ATTCTTGCCCT TGTACCTCAA TCTATTCTGT AAAATTAAAC AAATATGCAA     1616
ACCAGGATTT CCTTGACTTC TTTG                                     1640

```

- (2) INFORMATION FOR SEQUENCE ID NO: 12:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 943 base pairs
    - (B) TYPE: nucleic acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: genomic DNA
  - (ix) FEATURE:
    - (A) NAME/KEY: MAGE-31 gene
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

```

GGATCCTCCA CCCCAGTAGA GTGGGGACCT CACAGAGTCT GGCCAACCCCT      50
CCTGACAGTT CTGGGAATCC GTGGCTGCGT TTGCTGTCTG CACATTGGGG      100
GCCCCGTGGAT TCCTCTCCA GGAATCAGGA GCTCCAGGAA CAAGGCAGTG      150
AGGACTTGGT CTGAGGCAGT GTCCTCAGGT CACAGAGTAG AGGGGgCTCA      200
GATAGTGCCA ACGGTGAAGG TTTGCCTTGG ATTCAAACCA AGGGCCCCAC      250
CTGCCCCAGA ACACATGGAC TCCAGAGCGC CTGGCCTCAC CCTCAATACT      300
TTCAGTCCTG CAGCCTCAGC ATGCGCTGGC CGGATGTACC CTGAGGTGCC      350
CTCTCACTTC CTCCTTCAGG TTCTGAGGGG ACAGGCTGAC CTGGAGGACC      400
AGAGGCCCCC GGAGGAGCAC TGAAGGAGAA GATCTGTAAG TAAGCCTTTG      450
TTAGAGCCTC CAAGGTTCCA TTCAGTACTC AGCTGAGGTC TCTCACATGC      500
TCCCTCTCTC CCCAGGCCAG TGGGTCTCCA TTGCCAGCT CCTGCCACA      550
CTCCCGCCTG TTGCCCTGAC CAGAGTCATC      580
ATG CCT CTT GAG CAG AGG AGT CAG CAC TGC AAG CCT GAA GAA      622
GGC CTT GAG GCC CGA GGA GAg GCC CTG GGC CTG GTG GGT GCG      664
CAG GCT CCT GCT ACT GAG GAG CAG GAG GCT GCC TCC TCC TCT      706
TCT AGT GTA GTT GAA GTC ACC CTG GGG GAG GTG CCT GCT GCC      748
GAG TCA CCA GAT CCT CCC CAG AGT CCT CAG GGA GCC TCC AGC      790
CTC CCC ACT ACC ATG AAC TAC CCT CTC TGG AGC CAA TCC TAT      832
GAG GAC TCC AGC AAC CAA GAA GAG GAG GGG CCA AGC ACC TTC      874
CCT GAC CTG GAG TCT GAG TTC CAA GCA GCA CTC AGT AGG AAG      916
GTG GCC AAG TTG GTT CAT TTT CTG CTC      943

```

- (2) INFORMATION FOR SEQUENCE ID NO: 13:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 2531 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: MAGE-4 gene  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

```

GGATCCAGGC CCTGCCTGGA GAAATGTGAG GGCCCTGAGT GAACACAGTG      50
GGGATCATCC ACTCCATGAG AGTGGGGACC TCACAGAGTC CAGCCTACCC      100
TCTTGATGGC ACTGAGGGAC CGGGGCTGTG CTTACAGTCT GCACCCTAAG      150
GGCCCATGGA TTCCTCTCCT AGGAGCTCCA GGAACAAGGC AGTGAGGCCT      200
TGGTCTGAGA CAGTGTCTCTC AGGTTACAGA GCAGAGGATG CACAGGCTGT      250
GCCAGCAGTG AATGTTTGCC CTGAATGCAC ACCAAGGGGC CCACCTGCCA      300
CAAGACACAT AGGACTCCAA AGAGTCTGGC CTCACCTCCC TACCATCAAT      350
CCTGCAGAAAT CGACCTCTGC TGGCCGGCTA TACCCTGAGG TGCTCTCTCA      400
CTTCCTCCTT CAGGTTCTGA GCAGACAGGC CAACCGGAGA CAGGATTCCC      450
TGGAGGCCAC AGAGGAGCAC CAAGGAGAAG ATCTGTAAGT AAGCCTTTGT      500
TAGAGCCTCT AAGATTTGGT TCTCAGCTGA GGTCTCTCAC ATGCTCCCTC      550
TCTCCGTAGG CCTGTGGGTC CCCATTGCCC AGCTTTTGCC TGCACCTTG      600
CCTGCTGCCC TGACCAGAGT CATC                                     624
ATG TCT TCT GAG CAG AAG AGT CAG CAC TGC AAG CCT GAG GAA      666
GGC GTT GAG GCC CAA GAA GAG GAG GAG GAG GCT GCT GTC TCC TCC      708
CAG GCT CCT ACT ACT GAG GAG CAG GAG GCT GCT GTC TCC TCC      750
TCC TCT CCT CTG GTC CCT GGC ACC CTG GAG GAA GTG CCT GCT      792
GCT GAG TCA GCA GGT CCT CCC CAG AGT CCT CAG GGA GCC TCT      834
GCC TTA CCC ACT ACC ATC AGC TTC ACT TGC TGG AGG CAA CCC      876
AAT GAG GGT TCC AGC AGC CAA GAA GAG GAG GGG CCA AGC ACC      918
TCG CCT GAC GCA GAG TCC TTG TTC CGA GAA GCA CTC AGT AAC      960
AAG GTG GAT GAG TTG GCT CAT TTT CTG CTC CGC AAG TAT CGA      1002
GCC AAG GAG CTG GTC ACA AAG GCA GAA ATG CTG GAG AGA GTC      1044
ATC AAA AAT TAC AAG CGC TGC TTT CCT GTG ATC TTC GGC AAA      1086
GCC TCC GAG TCC CTG AAG ATG ATC TTT GGC ATT GAC GTG AAG      1128
GAA GTG GAC CCC GCC AGC AAC ACC TAC ACC CTT GTC ACC TGC      1170
CTG GGC CTT TCC TAT GAT GGC CTG CTG GGT AAT AAT CAG ATC      1212
TTT CCC AAG ACA GGC CTT CTG ATA ATC GTC CTG GGC ACA ATT      1254
GCA ATG GAG GGC GAC AGC GCC TCT GAG GAG GAA ATC TGG GAG      1296
GAG CTG GGT GTG ATG GGG GTG TAT GAT GGG AGG GAG CAC ACT      1338
GTC TAT GGG GAG CCC AGG AAA CTG CTC ACC CAA GAT TGG GTG      1380
CAG GAA AAC TAC CTG GAG TAC CGG CAG GTA CCC GGC AGT AAT      1422
CCT GCG CGC TAT GAG TTC CTG TGG GGT CCA AGG GCT CTG GCT      1464
GAA ACC AGC TAT GTG AAA GTC CTG GAG CAT GTG GTC AGG GTC      1506
AAT GCA AGA GTT CGC ATT GCC TAC CCA TCC CTG CGT GAA GCA      1548
GCT TTG TTA GAG GAG GAA GAG GGA GTC TGA                        1578
GCATGAGTTG CAGCCAGGGC TGTGGGGAAG GGGCAGGGCT GGGCCAGTGC      1628
ATCTAACAGC CCTGTGCAGC AGCTTCCCTT GCCTCGTGTA ACATGAGGCC      1678
CATTCTTCAC TCTGTTTGAA GAAAATAGTC AGTGTCTCTA GTAGTGGGTT      1728
TCTATTTTGT TGGATGACTT GGAGATTTAT CTCTGTTTCC TTTTACAATT      1778
GTTGAAATGT TCCTTTTAAT GGATGGTTGA ATTAACCTCA GCATCCAAGT      1828
TTATGAATCG TAGTTAACGT ATATTGCTGT TAATATAGTT TAGGAGTAAG      1878
AGTCTTGTTT TTTATTCAGA TTGGGAAATC CGTTCTATTT TGTGAATTTG      1928
GGACATAATA ACAGCAGTGG AGTAAGTATT TAGAAGTGTG AATTCACCGT      1978
GAAATAGGTG AGATAAATTA AAAGATACTT AATTCCTGCC TTATGCCTCA      2028
GTCTATTCTG TAAAATTTAA AAATATATAT GCATACCTGG ATTTCCCTTG      2078
CTTCGTGAAT GTAAGAGAAA TTAAATCTGA ATAAATAATT CTTTCTGTGA      2128
ACTGGCTCAT TTCTTCTCTA TGCACTGAGC ATCTGCTCTG TGGAAGGCCC      2178
AGGATTAGTA GTGGAGATAC TAGGGTAAGC CAGACACACA CCTACCGATA      2228
GGGTATTAGG AGTCTAGGAG CGCGGTCTAT TAATTAAGGT GACAAGATGT      2278
CCTCTAAGAT GTAGGGGAAA AGTAACGAGT GTGGGTATGG GGCTCCAGGT      2328
GAGAGTGGTC GGGTGTAAT TCCCTGTGTG GGGCCTTTTG GGCTTTGGGA      2378
AACTGCATTT TCTTCTGAGG GATCTGATTC TAATGAAGCT TGGTGGGTCC      2428
AGGGCCAGAT TCTCAGAGGG AGAGGGAAAA GCCCAGATTG GAAAAGTTGC      2478

```

67

TCTGAGCAGT TCCTTTGTGA CAATGGATGA ACAGAGAGGA GCCTCTACCT  
GGG

2528  
2531

- (2) INFORMATION FOR SEQUENCE ID NO: 14:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 2531 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: MAGE-41 gene  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

```

GGATCCAGGC CCTGCCTGGA GAAATGTGAG GGCCCTGAGT GAACACAGTG      50
GGGATCATCC ACTCCATGAG AGTGGGGACC TCACAGAGTC CAGCCTACCC      100
TCTTGATGGC ACTGAGGGAC CGGGGCTGTG CTTACAGTCT GCACCCTAAG      150
GGCCCATGGA TTCCTCTCCT AGGAGCTCCA GGAACAAGGC AGTGAGGCCT      200
TGGTCTGAGA CAGTGTCTCT AGGTTACAGA GCAGAGGATG CACAGGCTGT      250
GCCAGCAGTG AATGTTTGCC CTGAATGCAC ACCAAGGGCC CCACCTGCCA      300
CAAGACACAT AGGACTCCAA AGAGTCTGGC CTCACCTCCC TACCATCAAT      350
CCTGCAGAAT CGACCTCTGC TGGCCGGCTA TACCCTGAGG TGCTCTCTCA      400
CTTCCTCCTT CAGGTTCTGA GCAGACAGGC CAACCGGAGA CAGGATTCCC      450
TGGAGGCCAC AGAGGAGCAC CAAGGAGAAG ATCTGTAAAG AAGCCTTTGT      500
TAGAGCCTCT AAGATTTGGT TCTCAGCTGA GGTCTCTCAC ATGCTCCCTC      550
TCTCCGTAGG CTTGTGGGTC CCCATTGCCC AGCTTTTGCC TGCACCTTTG      600
CCTGCTGCCC TGAGCAGAGT CATC                                     624
ATG TCT TCT GAG CAG AAG AGT CAG CAC TGC AAG CCT GAG GAA      666
GGC GTT GAG GCC CAA GAA GAG GCC CTG GGC CTG GTG GGT GCG      708
CAG GCT CCT ACT ACT GAG GAG CAG GAG GCT GCT GTC TCC TCC      750
TCC TCT CCT CTG GTC CCT GGC ACC CTG GAG GAA GTG CCT GCT      792
GCT GAG TCA GCA GGT CCT CCC CAG AGT CCT CAG GGA GCC TCT      834
GCC TTA CCC ACT ACC ATC AGC TTC ACT TGC TGG AGG CAA CCC      876
AAT GAG GGT TCC AGC AGC CAA GAA GAG GAG GGG CCA AGC ACC      918
TCG CCT GAC GCA GAG TCC TTG TTC CGA GAA GCA CTC AGT AAC      960
AAG GTG GAT GAG TTG GCT CAT TTT CTG CTC CGC AAG TAT CGA      1002
GCC AAG GAG CTG GTC ACA AAG GCA GAA ATG CTG GAG AGA GTC      1044
ATC AAA AAT TAC AAG CGC TGC TTT CCT GTG ATC TTC GGC AAA      1086
GCC TCC GAG TCC CTG AAG ATG ATC TTT GGC ATT GAC GTG AAG      1128
GAA GTG GAC CCC ACC AGC AAC ACC TAC ACC CTT GTC ACC TGC      1170
CTG GGC CTT TCC TAT GAT GGC CTG CTG GGT AAT AAT CAG ATC      1212
TTT CCC AAG ACA GGC CTT CTG ATA ATC GTC CTG GGC ACA ATT      1254
GCA ATG GAG GGC GAC AGC GCC TCT GAG GAG GAA ATC TGG GAG      1296
GAG CTG GGT GTG ATG GGG GTG TAT GAT GGG AGG GAG CAC ACT      1338
GTC TAT GGG GAG CCC AGG AAA CTG CTC ACC CAA GAT TGG GTG      1380
CAG GAA AAC TAC CTG GAG TAC CGG CAG GTA CCC GGC AGT AAT      1422
CCT GCG CGC TAT GAG TTC CTG TGG GGT CCA AGG GCT CTG GCT      1464
GAA ACC AGC TAT GTG AAA GTC CTG GAG CAT GTG GTC AGG GTC      1506
AAT GCA AGA GTT CGC ATT GCC TAC CCA TCC CTG CGT GAA GCA      1548
GCT TTG TTA GAG GAG GAA GAG GGA GTC TGA                        1578
GCATGAGTTG CAGCCAGGGC TGTGGGGAAG GGGCAGGGCT GGGCCAGTGC      1628
ATCTAACAGC CCTGTGCAGC AGCTTCCCTT GCCTCGTGTA ACATGAGGCC      1678
CATCTTTCAC TCTGTTTGAA GAAAATAGTC AGTGTTCTTA GTAGTGGGTT      1728
TCTATTTTGT TGGATGACTT GGAGATTTAT CTCTGTTTCC TTTTACAATT      1778
GTTGAAATGT TCCTTTTAAAT GGATGGTTGA ATTAACCTCA GCATCCAAGT      1828
TTATGAATCG TAGTTAACGT ATATTGCTGT TAATATAGTT TAGGAGTAAG      1878
AGTCTTGTTT TTTATTCAGA TTGGGAAATC CGTTCTATTT TGTGAATTTG      1928
GGACATAATA ACAGCAGTGG AGTAAGTATT TAGAAGTGTG AATTCACCGT      1978
GAAATAGGTG AGATAAATTA AAAGATACTT AATTCCCGCC TTATGCCTCA      2028
GTCTATTCTG TAAAATTTAA AAATATATAT GCATACCTGG ATTTCCCTTG      2078
CTTCTGTAAT GTAAGAGAAA TTAAATCTGA ATAAATAATT CTTTCTGTTA      2128
ACTGGCTCAT TTCTTCTCTA TGCACCTGAGC ATCTGCTCTG TGGGAAGGCC      2178
AGGATTAGTA GTGGAGATAC TAGGGTAAGC CAGACACACA CCTACCGATA      2228
GGGTATTAAG AGTCTAGGAG CGCGGTCATA TAATTAAGGT GACAAGATGT      2278
CCTCTAAGAT GTAGGGGAAA AGTAACGAGT GTGGGTATGG GGCTCCAGGT      2328
GAGAGTGGTC GGGTGTAAAT TCCCTGTGTG GGGCCTTTTG GGCTTTGGGA      2378
AACTCCATTT TCTTCTGAGG GATCTGATTG TAATGAAGCT TGGTGGGTCC      2428

```

AGGGCCAGAT	TCTCAGAGGG	AGAGGGAAAA	GCCCAGATTG	GAAAAGTTGC	2478
TCTGAGCGGT	TCCTTTGTGA	CAATGGATGA	ACAGAGAGGA	GCCTCTACCT	2528
GGG					2531

- (2) INFORMATION FOR SEQUENCE ID NO: 15:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 1068 base pairs
    - (B) TYPE: nucleic acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: cDNA to mRNA
  - (ix) FEATURE:
    - (A) NAME/KEY: cDNA MAGE-4
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

G GGG CCA AGC ACC TCG CCT GAC GCA GAG TCC TTG TTC CGA	40
GAA GCA CTC AGT AAC AAG GTG GAT GAG TTG GCT CAT TTT CTG	82
CTC CGC AAG TAT CGA GCC AAG GAG CTG GTC ACA AAG GCA GAA	124
ATG CTG GAG AGA GTC ATC AAA AAT TAC AAG CGC TGC TTT CCT	166
GTG ATC TTC GGC AAA GCC TCC GAG TCC CTG AAG ATG ATC TTT	208
GGC ATT GAC GTG AAG GAA GTG GAC CCC GCC AGC AAC ACC TAC	250
ACC CTT GTC ACC TGC CTG GGC CTT TCC TAT GAT GGC CTG CTG	292
GGT AAT AAT CAG ATC TTT CCC AAG ACA GGC CTT CTG ATA ATC	334
GTC CTG GGC ACA ATT GCA ATG GAG GGC GAC AGC GCC TCT GAG	376
GAG GAA ATC TGG GAG GAG CTG GGT GTG ATG GGG GTG TAT GAT	418
GGG AGG GAG CAC ACT GTC TAT GGG GAG CCC AGG AAA CTG CTC	460
ACC CAA GAT TGG GTG CAG GAA AAC TAC CTG GAG TAC CGG CAG	502
GTA CCC GGC AGT AAT CCT GCG CGC TAT GAG TTC CTG TGG GGT	544
CCA AGG GCT CTG GCT GAA ACC AGC TAT GTG AAA GTC CTG GAG	586
CAT GTG GTC AGG GTC AAT GCA AGA GTT CGC ATT GCC TAC CCA	628
TCC CTG CGT GAA GCA GCT TTG TTA GAG GAG GAA GAG GGA GTC	670
TGAGCATGAG TTGCAGCCAG GGCTGTGGGG AAGGGGCAGG GCTGGGCCAG	720
TGCATCTAAC AGCCCTGTGC AGCAGCTTCC CTTGCCTCGT GTAACATGAG	770
GCCCATTCTT CACTCTGTTT GAAGAAAATA GTCAGTGTTT TTAGTAGTGG	820
GTTTCTATTT TGTTGGATGA CTTGGAGATT TATCTCTGTT TCCTTTTACA	870
ATTGTTGAAA TGTTCTTTT AATGGATGGT TGAATTAAC TCAGCATCCA	920
AGTTTATGAA TCGTAGTTAA CGTATATTGC TGTTAATATA GTTTAGGAGT	970
AAGAGTCTTG TTTTATTTC AGATTGGGAA ATCCGTTCTA TTTTGTGAAT	1020
TTGGGACATA ATAACAGCAG TGGAGTAAGT ATTTAGAAGT GTGAATTC	1068



- (2) INFORMATION FOR SEQUENCE ID NO: 16:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 2226 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: MAGE-5 gene  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

GGATCCAGGC	CTTGCCAGGA	GAAAGGTGAG	GGCCCTGTGT	GAGCACAGAG	50
GGGACCATT	ACCCCAAGAG	GGTGGAGACC	TCACAGATT	CAGCCTACCC	100
TCCTGTTAG	ACTGGGGGCC	TGAGGCTGTG	CTTGCACTCT	GCACCCCTGAG	150
GGCCCATGCA	TTCCTCTTCC	AGGAGCTCCA	GGAAACAGAC	ACTGAGGCCT	200
TGGTCTGAG	CCGTGCCCTC	AGGTCACAGA	GCAGAGGAGA	TGCAGACGTC	250
TAGTGCCAG	AGTGAACGTT	TGCCTTGAAT	GCACACTAAT	GGCCCCCATC	300
GCCCCAGAAC	ATATGGGACT	CCAGAGCACC	TGGCCTCACC	CTCTCTACTG	350
TCAGTCCTGC	AGAATCAGCC	TCTGCTTGCT	TGTGTACCCT	GAGGTGCCCT	400
CTCACTTTTT	CCTTCAGGTT	CTCAGGGGAC	AGGCTGACCA	GGATCACCAG	450
GAAGCTCCAG	AGGATCCCCA	GGAGGCCCTA	GAGGAGCACC	AAAGGAGAAG	500
ATCTGTAAGT	AAGCCTTTGT	TAGAGCCTCC	AAGGTTCACT	TTTGTAGCTGA	550
GGCTTCTCAC	ATGCTCCCTC	TCTCTCCAGG	CCAGTGGGTC	TCCATTGCCC	600
AGCTCCTGCC	CACACTCCTG	CCTGTTGCGG	TGACCAGAGT	CGTC	644
ATG TCT CTT	GAG CAG AAG	AGT CAG CAC	TGC AAG CCT	GAG GAA	684
CTC CTC TGG	TCC CAG GCA	CCC TGG GGG	AGG TGC CTG	CTG CTG CTG	728
GGT CAC CAG	GTC CTC TCA	AGA GTC CTC	AGG GAG CCT	CCG CCA	770
TCC CCA CTG	CCA TCG ATT	TCA CTC TAT	GGA GGC AAT	CCA TTA	812
AGG GCT CCA	GCA ACC AAG	AAG AGG AGG	GGC CAA GCA	CCT CCC	854
CTG ACC CAG	AGT CTG TGT	TCC GAG CAG	CAC TCA GTA	AGA AGG	896
TGG CTG ACT	TGA				908
TTCATTTTTCT	GCTCCTCAAG	TATTAAGTCA	AGGAGCTGGT	CACAAAGGCA	958
GAAATGCTGG	AGAGCGTCAT	CAAAAATTAC	AAGCGCTGCT	TTCCTGAGAT	1008
CTTCGGCAAA	GCCTCCGAGT	CCTTGCAGCT	GGTCTTTGGC	ATTGACGTGA	1058
AGGAAGCGGA	CCCCACCAGC	AACACCTACA	CCCTTGTCA	CTGCCTGGGA	1108
CTCCTATGAT	GGCCTGCTGG	TTGATAATAA	TCAGATCATG	CCCAAGACGG	1158
GCCTCCTGAT	AATCGTCTTG	GGCATGATTG	CAATGGAGGG	CAAAATGCGTC	1208
CCTGAGGAGA	AAATCTGGGA	GGAGCTGAGT	GTGATGAAGG	TGTATGTTGG	1258
GAGGGAGCAC	AGTGTCTGTG	GGGAGCCCAG	GAAGCTGCTC	ACCCAAGATT	1308
TGGTGCAGGA	AAACTACCTG	GAGTACCGGC	AGGTGCCAG	CAGTGATCCC	1358
ATATGCTATG	AGTTACTGTG	GGGTCCAAGG	GCACTCGCTG	CTTGAAAGTA	1408
CTGGAGCACG	TGGTCAGGGT	CAATGCAAGA	GTTCTCATTT	CCTACCCATC	1458
CCTGCGTGAA	GCAGCTTTGA	GAGAGGAGGA	AGAGGGAGTC	TGAGCATGAG	1508
CTGCAGCCAG	GGCCAATGCG	AGGGGGGCTG	GGCCAGTGCA	CCTTCCAGGG	1558
CTCCGTCCAG	TAGTTTCCCC	TGCCTTAATG	TGACATGAGG	CCCATTCTTC	1608
TCTCTTTGAA	GAGAGCAGTC	AACATTCTTA	GTAGTGGGTT	TCTGTTCTAT	1658
TGGATGACTT	TGAGATTGTG	CTTTGTTTCC	TTTTGGAATT	GTTCAAATGT	1708
TTCTTTTAAT	GGGTGGTTGA	ATGAACTTCA	GCATTCAAAT	TTATGAATGA	1758
CAGTAGTCAC	ACATAGTGCT	GTTTATATAG	TTTAGGAGTA	AGAGTCTTGT	1808
TTTTTATTCA	GATTGGGAAA	TCCATTCCAT	TTTGTGAATT	GGGACATAGT	1858
TACAGCAGTG	GAATAAGTAT	TCATTTAGAA	ATGTGAATGA	GCAGTAAAAC	1908
TGATGACATA	AAGAAATTAA	AAGATATTTA	ATTCTTGCTT	ATACTCAGTC	1958
TATTCGGTAA	AATTTTTTTT	AAAAAATGTG	CATACCTGGA	TTTCCTTGGC	2008
TTCTTTGAGA	ATGTAAGACA	AATTAAATCT	GAATAAATCA	TTCTCCCTGT	2058
TCACTGGGCTC	ATTTATTCTC	TATGCACTGA	GCATTGCTC	TGTGGAAGGC	2108
CCTGGGTTAA	TAGTGAGAT	GCTAAGGTAA	GCCAGACTCA	CCCCTACCCA	2158
CAGGGTAGTA	AAGTCTAGGA	GCAGCAGTCA	TATAATTAAG	GTGAGAGAT	2208
GCCCTCTAAG	ATGTAGAG				2226

- (2) INFORMATION FOR SEQUENCE ID NO: 17:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 2305 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: MAGE-51 gene  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

```

GGATCCAGGC CTTGCCAGGA GAAAGGTGAG GGCCCTGTGT GAGCACAGAG      50
GGGACCATTG ACCCAAGAG GGTGGAGACC TCACAGATTG CAGCCTACCC      100
TCCTGTTAGC ACTGGGGGCC TGAGGCTGTG CTTGCAGTCT GCACCCCTGAG      150
GGCCCATGCA TTCCTCTTCC AGGAGCTCCA GGAAACAGAC ACTGAGGCCT      200
TGGTCTGAGG CCGTGCCCTC AGGTACAGAG GCAGAGGAGA TGCAGACGTC      250
TAGTGCCAGC AGTGAACGTT TGCCTTGAAT GCACCTAAT GGCCCCCATC      300
GCCCCAGAAC ATATGGGACT CCAGAGCACC TGGCCTCACC CTCTCTACTG      350
TCAGTCCTGC AGAATCAGCC TCTGCTTGCT TGTGTACCCT GAGGTGCCCT      400
CTCACTTTTT CCTTCAGGTT CTCAGGGGAC AGGCTGACCA GGATCACCAG      450
GAAGCTCCAG AGGATCCCCA GGAGGCCCTA GAGGAGCACC AAAGGAGAAG      500
ATCTGTAAGT AAGCCTTTGT TAGAGCCTCC AAGGTTTCACT TTTAGCTGA      550
GGCTTCTCAC ATGCTCCCTC TCTCTCCAGG CCAGTGGGTC TCCATTGCCC      600
AGCTCCTGCC CACACTCCTG CCTGTTGCGG TGACCAGAGT CGTC          644
ATG TCT CTT GAG CAG AAG AGT CAG CAC TGC AAG CCT GAG GAA      686
GGC CTT GAC ACC CAA GAA GAG CCC TGG GCC TGG TGG GTG TGC      728
AGG CTG CCA CTA CTG AGG AGC AGG AGG CTG TGT CCT CCT CCT      770
CTC CTC TGG TCC CAG GCA CCC TGG GGG AGG TGC CTG CTG CTG      812
GGT CAC CAG GTC CTC TCA AGA GTC CTC AGG GAG CCT CCG CCA      854
TCC CCA CTG CCA TCG ATT TCA CTC TAT GGA GGC AAT CCA TTA      896
AGG GCT CCA GCA ACC AAG AAG AGG AGG GGC CAA GCA CCT CCC      938
CTG ACC CAG AGT CTG TGT TCC GAG CAG CAC TCA GTA AGA AGG      980
TGG CTG ACT TGA
TTCATTTTCT GCTCCTCAAG TATTAAGTCA AGGAGCCGGT CACAAAGGCA      1042
GAAATGCTGG AGAGCGTCAT CAAAAATTAC AAGCGCTGCT TTCCTGAGAT      1092
CTTCGGCAAA GCCTCCGAGT CCTTGCAGCT GGTCTTTGGC ATTGACGTGA      1142
AGGAAGCGGA CCCCACCAGC AACACCTACA CCCTTGTAC CTGCTGGGA      1192
CTCCTATGAT GGCCTGGTGG TTTAATCAGA TCATGCCCAA GACGGGCCTC      1242
CTGATAATCG TCTTGGGCAT GATTGCAATG GAGGGCAAAT GCGTCCCTGA      1292
GGAGAAAATC TGGGAGGAGC TGGGTGTGAT GAAGGTGTAT GTTGGGAGGG      1342
AGCACAGTGT CTGTGGGGAG CCCAGGAAGC TGCTCACCCA AGATTGGTG      1392
CAGGAAAATC ACCTGGAGTA CCGCAGGTGC CCAGCAGTGA TCCCATATGC      1442
TATGAGTTAC TGTGGGGTCC AAGGGCACTC GCTGCTTGAA AGTACTGGAG      1492
CACGTGGTCA GGGTCAATGC AAGAGTTCTC ATTTCTACC CATCCCTGCA      1542
TGAAGCAGCT TTGAGAGAGG AGGAAGAGGG AGTCTGAGCA TGAGCTGCAG      1592
CCAGGGCCAC TGCGAGGGGG GCTGGGCCAG TGCACCTTCC AGGGCTCCGT      1642
CCAGTAGTTT CCCCTGCCTT AATGTGACAT GAGGCCATT CTCTCTCTT      1692
TGAAGAGAGC AGTCAACATT CTTAGTAGTG GGTTCCTGTT CTATTGGATG      1742
ACTTTGAGAT TTGTCTTTGT TTCCTTTTGG AATTGTTCAA ATGTTCTTTT      1792
TAATGGGTGG TTGAATGAAC TTCAGCATTG AAATTTATGA ATGACAGTAG      1842
TCACACATAG TGCTGTTTAT ATAGTTTAGG AGTAAGAGTC TTGTTTTTTA      1892
TTCAGATTGG GAAATCCATT CCATTTTGTG AATTGGGACA TAGTTACAGC      1942
AGTGGAATAA GTATTCATT AGAAATGTGA ATGAGCAGTA AAACCTGATGA      1992
GATAAAGAAA TTAAAGATA TTTAATTCTT GCCTTATACT CAGTCTATTC      2042
GGTAAATTTT TTTTTTAAA ATGTGCATAC CTGGATTTCC TTGGCTTCTT      2092
TGAGAAATGA AGACAAATTA AATCTGAATA AATCATTCTC CCTGTTCACT      2142
GGCTCATTTA TTCTCTATGC ACTGAGCATT TGCTCTGTGG AAGGCCCTGG      2192
GTTAATAGTG GAGATGCTAA GGTAAGCCAG ACTCACCCCT ACCCACAGGG      2242
TAGTAAAGTC TAGGAGCAGC AGTCATATAA TTAAGGTGGA GAGATGCCCT      2292
CTAAGATGTA GAG

```

- (2) INFORMATION FOR SEQUENCE ID NO: 18:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 225 base pairs
    - (B) TYPE: nucleic acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: cDNA
  - (ix) FEATURE:
    - (A) NAME/KEY: MAGE-6 gene
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

TAT TTC TTT CCT GTG ATC TTC AGC AAA GCT TCC GAT TCC TTG	42
CAG CTG GTC TTT GGC ATC GAG CTG ATG GAA GTG GAC CCC ATC	84
GGC CAC GTG TAC ATC TTT GCC ACC TGC CTG GGC CTC TCC TAC	126
GAT GGC CTG CTG GGT GAC AAT CAG ATC ATG CCC AGG ACA GGC	168
TTC CTG ATA ATC ATC CTG GCC ATA ATC GCA AGA GAG GGC GAC	210
TGT GCC CCT GAG GAG	225

- (2) INFORMATION FOR SEQUENCE ID NO: 19:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 1947 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: MAGE-7 gene  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

TGAATGGACA	ACAAGGGCCC	CACACTCCCC	AGAACACAAG	GGA	50
GAGCCCAGCC	TCACCTTCCC	TACTGTTCAGT	CCTGCAAGCCT	CAGCCTCTGC	100
TGGCCGGCTG	TACCTTGAGG	TGCCCTCTCA	CTTCCCTCCTT	CAGGTTCTCA	150
GCGGACAGGC	CGGCCAGGAG	GTCAGAAAGCC	CCAGGAGGCC	CCAGAGGAGC	200
ACCGAAGGAG	AAGATCTGTA	AGTAGGCCTT	TGTTAGGGCC	TCCAGGGCGT	250
GGTTCACAAA	TGAGGCCCCCT	CACAAGCTCC	TTCTCTCCCC	AGATCTGTGG	300
GTTCCTCCCC	ATCGCCCGAGC	TGCTGCCCGC	ACTCAGCCT	GCTGCCCTGA	350
CCAGAGTCAT	CATGTCTTCT	GAGCAGAGGA	GTCAGCACTG	CAAGCCTGAG	400
GATGCCTTGA	GGCCCAAGGA	CAGGAGGCTC	TGGGCCTGGT	GGGTGCGCAG	450
GCTCCCGCCA	CCGAGGAGCA	CGAGGCTGCC	TCCTCCTTCA	CTCTGATTGA	500
AGGCACCCCT	GAGGAGGTGC	CTGCTGCTGG	GTCCCCCAGT	CCTCCCTGA	550
GTCTCAGGGT	TCCTCCTTTT	CCCTGACCAT	CAGCAACAAC	ACTCTATGGA	600
GCCAATCCAG	TGAGGGCACC	AGCAGCCGGG	AAGAGGAGGG	GCCAACCACC	650
TAGACACACC	CCGCTCACCT	GGCGTCCTTG	TTCCA		685
ATG GGA AGG	TGG CTG AGT	TGG TTC GCT	TCC TGC	TGC ACA AGT	727
ATC GAG TCA	AGG AGC TGG	TCA CAA AGG	CAG AAA TGC	TGG ACA	769
GTG TCA TCA	AAA ATT ACA	AGC ACT AGT	TTC CTT GTG	ATC TAT	811
GGC AAA GCC	TCA GAG TGC	ATG CAG GTG	ATG TTT GGC	ATT GAC	853
ATG AAG GAA	GTG GAC CCC	GCG GCC ACT	CCT ACG TCC	TTG TCA	895
CCT GCT TGG	GCC TCT CCT	ACA ATG GCC	TGC TGG GTG	ATG ATC	937
AGA GCA TGC	CCG AGA CCG	GCC TTC TGA			964
TTATGGTCTT	GACCATGATC	TTAATGGAGG	GCCACTGTGC	CCCTGAGGAG	1014
GCAATCTGGG	AAGCGTTGAG	TGTAATGGTG	TATGATGGGA	TGGAGCAGTT	1064
TCTTTGGGCA	GCTGAGGAAG	CTGCTCACCC	AAGATTGGGT	GCAGGAAAAC	1114
TACCTGCAAT	ACCGCCAGGT	GCCCAGCAGT	GATCCCCCGT	GCTACCAAGT	1164
CCTGTGGGGT	CCAAGGGCCC	TCATTGAAAC	CAGCTATGTG	AAAGTCCTGG	1214
AGTATGCAGC	CAGGGTCAGT	ACTAAAGAGA	GCATTTCTTA	CCCATCCCTG	1264
CATGAAGAGG	CTTTGGGAGA	GGAGGAAGAG	GGAGTCTGAG	CAGAAGTTGC	1314
AGCCAGGGCC	AGTGGGGCAG	ATTGGGGGAG	GGCCTGGGCA	GTGCACGTTC	1364
CACACATCCA	CCACCTTCCC	TGTCTGTGTA	CATGAGGCCC	ATTCTTCACT	1414
CTGTGTTTGA	AGAGAGCAGT	CAATGTTCTC	AGTAGCGGGG	AGTGTGTTGG	1464
GTGTGAGGGA	ATACAAGGTG	GACCATCTCT	CAGTTCCTGT	TCTCTTGGGC	1514
GATTTGGAGG	TTTATCTTTG	TTTCCTTTTG	CAGTCGTTCA	AATGTTCTTT	1564
TTAATGGATG	GTGTAATGAA	CTTCAACATT	CATTTCATGT	ATGACAGTAG	1614
GCAGACTTAC	TGTTTTTTAT	ATAGTTAAAA	GTAAGTGCAT	TGTTTTTTAT	1664
TTATGTAAGA	AAATCTATGT	TATTTCTTGA	ATTGGGACAA	CATAACATAG	1714
CAGAGGATTA	AGTACCTTTT	ATAATGTGAA	AGAACAAAGC	GGTAAAATGG	1764
GTGAGATAAA	GAAATAAAGA	AATTAAATTG	GCTGGGCACG	GTGGCTCACC	1814
CCTGTAATCC	CAGCACTTTA	GGAGGCAGAG	GCACGGGGAT	CACGAGGTCA	1864
GGAGATCGAG	ACCATTCTGG	CTAACACAGT	GAAACACCAT	CTCTATTAAA	1914
AATACAAAAC	TTAGCCGGGC	GTGGTGGCGG	GTG		1947

- (2) INFORMATION FOR SEQUENCE ID NO: 20:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 1810 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: MAGE-8 gene  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

```

GAGCTCCAGG AACCAGGCTG TGAGGTCTTG GTCTGAGGCA GTATCTTCAA      50
TCACAGAGCA TAAGAGGCCC AGGCAGTAGT AGCAGTCAAG CTGAGGTGGT      100
GTTTCCCCCTG TATGTATACC AGAGGCCCTT CTGGCATCAG AACAGCAGGA      150
ACCCACAGT TCCTGGCCCT ACCAGCCCTT TTGTCACTCC TGGAGCCTTG      200
GCCTTTGCCA GGAGGCTGCA CCCTGAGATG CCCTCTCAAT TTCTCCTTCA      250
GGTTCGCAGA GAACAGGCCA GCCAGGAGGT CAGGAGGCC CAGAGAAGCA      300
CTGAAGAAGA CCTGTAAGTA GACCTTTGTT AGGGCATCCA GGGTGTAGTA      350
CCCAGCTGAG GCCTCTCACA CGCTTCCTCT CTCCCAGGC CTGTGGGTCT      400
CAATTGCCCA GCTCCGGCCC AACTCTCCT GCTGCCCTGA CCTGAGTCAT      450
C
ATG CTT CTT GGG CAG AAG AGT CAG CGC TAC AAG GCT GAG GAA      493
GGC CTT CAG GCC CAA GGA GAG GCA CCA GGG CTT ATG GAT GTG      535
CAG ATT CCC ACA GCT GAG GAG CAG AAG GCT GCA TCC TCC TCC      577
TCT ACT CTG ATC ATG GGA ACC CTT GAG GAG GTG ACT GAT TCT      619
GGG TCA CCA AGT CCT CCC CAG AGT CCT GAG GGT GCC TCC TCT      661
TCC CTG ACT GTC ACC GAC AGC ACT CTG TGG AGC CAA TCC GAT      703
GAG GGT TCC AGC AGC AAT GAA GAG GAG GGG CCA AGC ACC TCC      745
CCG GAC CCA GCT CAC CTG GAG TCC CTG TTC CGG GAA GCA CTT      787
GAT GAG AAA GTG GCT GAG TTA GTT CGT TTC CTG CTC CGC AAA      829
TAT CAA ATT AAG GAG CCG GTC ACA AAG GCA GAA ATG CTT GAG      871
AGT GTC ATC AAA AAT TAC AAG AAC CAC TTT CCT GAT ATC TTC      913
AGC AAA GCC TCT GAG TGC ATG CAG GTG ATC TTT GGC ATT GAT      955
GTG AAG GAA GTG GAC CCT GCC GGC CAC TCC TAC ATC CTT GTC      997
ACC TGC CTG GGC CTC TCC TAT GAT GGC CTG CTG GGT GAT GAT      1039
CAG AGT ACG CCC AAG ACC GGC CTC CTG ATA ATC GTC CTG GGC      1081
ATG ATC TTA ATG GAG GGC AGC CGC GCC CCG GAG GAG GCA ATC      1123
TGG GAA GCA TTG AGT GTG ATG GGG GCT GTA TGA      1156
TGGGAGGGAG CACAGTGTCT ATTGGAAGCT CAGGAAGCTG CTCACCCAAG      1206
AGTGGGTGCA GGAGAACTAC CTGGAGTACC GCCAGGCGCC CGGCAGTGAT      1256
CCTGTGCGCT ACGAGTTCCT GTGGGGTCCA AGGGCCCTTG CTGAAACCAG      1306
CTATGTGAAA GTCCTGGAGC ATGTGGTCAG GGTCAATGCA AGAGTTCGCA      1356
TTTCCTACCC ATCCCTGCAT GAAGAGGCTT TGGGAGAGGA GAAAGGAGTT      1406
TGAGCAGGAG TTGCAGCTAG GGCCAGTGGG GCAGGTTGTG GGAGGGCCTG      1456
GGCCAGTGCA CGTTCCAGGG CCACATCCAC CACTTTCCTT GCTCTGTTAC      1506
ATGAGGCCCA TTCTTCACTC TGTGTTTGAA GAGAGCAGTC ACAGTTCTCA      1556
GTAGTGGGGA GCATGTTGGG TGTGAGGGAA CACAGTGTGG ACCATCTCTC      1606
AGTTCCTGTT CTATTGGGCG ATTTGGAGGT TTATCTTTGT TTCCTTTTGG      1656
AATTGTTCCA ATGTTTCCTT TAATGGATGG TGTAATGAAC TTCAACATTC      1706
ATTTTATGTA TGACAGTAGA CAGACTTACT GCTTTTTATA TAGTTTAGGA      1756
GTAAGAGTCT TGCTTTTCAT TTATACTGGG AAACCCATGT TATTTCTTGA      1806
ATTG      1810

```

- (2) INFORMATION FOR SEQUENCE ID NO: 21:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 1412 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: MAGE-9 gene  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

```

TCTGAGACAG TGTCCTCAGG TCGCAGAGCA GAGGAGACCC AGGCAGTGTC      50
AGCAGTGAAG GTGAAGTGTT CACCCTGAAT GTGCACCAAG GGCCCCACCT      100
GCCCCAGCAC ACATGGGACC CCATAGCACC TGGCCCCATT CCCCCTACTG      150
TCACTCATAG AGCCTTGATC TCTGCAGGCT AGCTGCACGC TGAGTAGCCC      200
TCTCACTTCC TCCCTCAGGT TCTCGGGACA GGCTAACCAG GAGGACAGGA      250
GCCCCAAGAG GCCCCAGAGC AGCACTGACG AAGACCTGTA AGTCAGCCTT      300
TGTTAGAACC TCCAAGGTTT GGTTCCTCAGC TGAAGTCTCT CACACACTCC      350
CTCTCTCCCC AGGCCTGTGG GTCTCCATCG CCCAGCTCCT GCCCAGCCTC      400
CTGACTGCTG CCCTGACCAG AGTCATC      427
ATG TCT CTC GAG CAG AGG AGT CCG CAC TGC AAG CCT GAT GAA      469
GAC CTT GAA GCC CAA GGA GAG GAC TTG GGC CTG ATG GGT GCA      511
CAG GAA CCC ACA GGC GAG GAG GAG GAG ACT ACC TCC TCC TCT      553
GAC AGC AAG GAG GAG GAG GTG TCT GCT GCT GGG TCA TCA AGT      595
CCT CCC CAG AGT CCT CAG GGA GGC GCT TCC TCC TCC ATT TCC      637
GTC TAC TAC ACT TTA TGG AGC CAA TTC GAT GAG GGC TCC AGC      679
AGT CAA GAA GAG GAA GAG CCA AGC TCC TCG GTC GAC CCA GCT      721
CAG CTG GAG TTC ATG TTC CAA GAA GCA CTG AAA TTG AAG GTG      763
GCT GAG TTG GTT CAT TTC CTG CTC CAC AAA TAT CGA GTC AAG      805
GAG CCG GTC ACA AAG GCA GAA ATG CTG GAG AGC GTC ATC AAA      847
AAT TAC AAG CGC TAC TTT CCT GTG ATC TTC GGC AAA GCC TCC      889
GAG TTC ATG CAG GTG ATC TTT GGC ACT GAT GTG AAG GAG GTG      931
GAC CCC GCC GGC CAC TCC TAC ATC CTT GTC ACT GCT CTT GGC      973
CTC TCG TGC GAT AGC ATG CTG GGT GAT GGT CAT AGC ATG CCC      1015
AAG GCC GCC CTC CTG ATC ATT GTC CTG GGT GTG ATC CTA ACC      1057
AAA GAC AAC TGC GCC CCT GAA GAG GTT ATC TGG GAA GCG TTG      1099
AGT GTG ATG GGG GTG TAT GTT GGG AAG GAG CAC ATG TTC TAC      1141
GGG GAG CCC AGG AAG CTG CTC ACC CAA GAT TGG GTG CAG GAA      1183
AAC TAC CTG GAG TAC CGG CAG GTG CCC GGC AGT GAT CCT GCG      1225
CAC TAC GAG TTC CTG TGG GGT TCC AAG GCC CAC GCT GAA ACC      1267
AGC TAT GAG AAG GTC ATA AAT TAT TTG GTC ATG CTC AAT GCA      1309
AGA GAG CCC ATC TGC TAC CCA TCC CTT TAT GAA GAG GTT TTG      1351
GGA GAG GAG CAA GAG GGA GTC TGA      1375
GCACCAGCCG CAGCCGGGGC CAAAGTTTGT GGGGTCA      1412

```

- (2) INFORMATION FOR SEQUENCE ID NO: 22:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 920 base pairs
    - (B) TYPE: nucleic acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: genomic DNA
  - (ix) FEATURE:
    - (A) NAME/KEY: MAGE-10 gene
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

ACCTGCTCCA	GGACAAAGTG	GACCCCACTG	CATCAGCTCC	ACCTACCCTA	50
CTGTCAGTCC	TGGAGCCTTG	GCCTCTGCCG	GCTGCATCCT	GAGGAGCCAT	100
CTCTCACTTC	CTTCTTCAGG	TTCTCAGGGG	ACAGGGAGAG	CAAGAGGTCA	150
AGAGCTGTGG	GACACCACAG	AGCAGCACTG	AAGGAGAAGA	CCTGTAAGTT	200
GGCCTTTGTT	AGAACCTCCA	GGGTGTGGTT	CTCAGCTGTG	GCCACTTACA	250
CCCTCCCTCT	CTCCCCAGGC	CTGTGGGTCC	CCATCGCCCA	AGTCCTGCCC	300
ACACTCCAC	CTGCTACCCT	GATCAGAGTC	ATC		333
ATG CCT CGA	GCT CCA AAG	CGT CAG CGC	TGC ATG CCT	GAA GAA	375
GAT CTT CAA	TCC CAA AGT	GAG ACA CAG	GGC CTC GAG	GGT GCA	417
CAG GCT CCC	CTG GCT GTG	GAG GAG GAT	GCT TCA TCA	TCC ACT	459
TCC ACC AGC	TCC TCT TTT	CCA TCC TCT	TTT CCC TCC	TCC TCC	501
TCT TCC TCC	TCC TCC TGC	TAT CCT CTA	ATA CCA AGC	ACC	543
CCA GAG GAG	GTT TCT GCT	GAT GAT GAG	ACA CCA AAT	CCT CCC	585
CAG AGT GCT	CAG ATA GCC	TGC TCC TCC	CCC TCG GTC	GTT GCT	627
TCC CTT CCA	TTA GAT CAA	TCT GAT GAG	GGC TCC AGC	AGC CAA	669
AAG GAG GAG	AGT CCA AGC	ACC CTA CAG	GTC CTG CCA	GAC AGT	711
GAG TCT TTA	CCC AGA AGT	GAG ATA GAT	GAA AAG GTG	ACT GAT	753
TTG GTG CAG	TTT CTG CTC	TTC AAG TAT	CAA ATG AAG	GAG CCG	795
ATC ACA AAG	GCA GAA ATA	CTG GAG AGT	GTC ATA AAA	AAT TAT	837
GAA GAC CAC	TTC CCT TTG	TTG TTT AGT	GAA GCC TCC	GAG TGC	879
ATG CTG CTG	GTC TTT GGC	ATT GAT GTA	AAG GAA GTG	GAT CC	920

(2) INFORMATION FOR SEQUENCE ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1107 base pairs

(B) TYPE: nucleic acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic DNA

(ix) FEATURE:

(A) NAME/KEY: MAGE-11 gene

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

AGAGAACAGG	CCAACTTGG	GGACAGGAGT	CCCAGGAGAA	CCCAGAGGAT	50
CACTGGAGGA	GAACAAGTGT	AAGTAGGCCT	TGTGTAGATT	CTCCATGGTT	100
CATATCTCAT	CTGAGTCTGT	TCTCAGCTC	CCTCTCTCCC	CAGGCTGTGG	150
GGCCCCATCA	CCGAGATATT	TCCACAGTT	CGCCTGCTG	ACCTAACCA	200
AGTCATCATG	CCTCTTGAGC	AAAGAAGTCA	GCACTGCAAG	CCTGAGGAAG	250
CCTTCAGGCC	CAAGAAGAAG	ACCTGGGCCT	GGTGGGTGCA	CAGGCTCTCC	300
AAGCTGAGGA	GCAGGAGGCT	GCCTTCTTCT	CCTCTACTCT	GAATGTGGGC	350
ACTCTAGAGG	AGTTGCCTGC	TGCTGAGTCA	CCAAGTCTCT	CCCAGAGTCC	400
TCAGGAAGAG	TCCTTCTCTC	CCACTGCCAT	GGATGCCATC	TTTGGGAGCC	450
TATCTGATGA	GGGCTCTGGC	AGCCAAGAAA	AGGAGGGGCC	AAGTACCTCG	500
CCTGACCTGA	TAGACCTTGA	TGCTCTTTCC	CAAGATATAC	TACATGACAA	550
GATAATTGAT	TTGGTTCATT	TATTCTCCGC	AAGTATCGAG	TCAAGGGGCT	600
GATCACAAAG	GCAGAA				616
ATG CTG GGG	AGT GTC ATC	AAA AAT TAT	GAG GAC TAC	TTT CCT	658
GAG ATA TTT	AGG GAA GCC	TCT GTA TGC	ATG CAA CTG	CTC TTT	700
GGC ATT GAT	GTG AAG GAA	GTG GAC CCC	ACT AGC CAC	TCC TAT	742
GTC CTT GTC	ACC TCC CTC	AAC CTC TCT	TAT GAT GGC	ATA CAG	784
TGT AAT GAG	CAG AGC ATG	CCC AAG TCT	GGC CTC CTG	ATA ATA	826
GTC CTG GGT	GTA ATC TTC	ATG GAG GGG	AAC TGC ATC	CCT GAA	868
GAG GTT ATG	TGG GAA GTC	CTG AGC ATT	ATG GGG GTG	TAT GCT	910
GGA AGG GAG	CAC TTC CTC	TTT GGG GAG	CCC AAG AGG	CTC CTT	952
ACC CAA AAT	TGG GTG CAG	GAA AAG TAC	CTG GTG TAC	CGG CAG	994
GTG CCC GGC	ACT GAT CCT	GCA TGC TAT	GAG TTC CTG	TGG GGT	1036
CCA AGG GCC	CAC GCT GAG	ACC AGC AAG	ATG AAA GTT	CTT GAG	1078
TAC ATA GCC	AAT GCC AAT	GGG AGG GAT	CC		1100



- (2) INFORMATION FOR SEQUENCE ID NO: 24:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 2150 base pairs
    - (B) TYPE: nucleic acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: genomic DNA
  - (ix) FEATURE:
    - (A) NAME/KEY: smage-I
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

TCTGCTCTGCA	TATGCCTCCA	CTTGTGTGTA	GCAGTCTCAA	ATGGATCTCT	50
CTCTACAGAC	CTCTGTCTGT	GTCTGGCACC	CTAAGTGGCT	TTGCATGGGC	100
ACAGGTTTCT	GGCCCTGCAT	GGAGCTTAAA	TAGATCTTTC	TCCACAGGCC	150
TATACCCCTG	CATTGTAAGT	TTAAGTGGCT	TTATGTGGAT	ACAGGTCTCT	200
GGCCTTGTAT	GCAGGCCTAA	GTTTTTCTGT	CTGCTTAACC	CCTCCAAGTG	250
AAGCTAGTGA	AAGATCTAAC	CCACTTTTGG	AAGTCTGAAA	CTAGACTTTT	300
ATGCAGTGGC	CTAACAAGTT	TTAATTTCTT	CCACAGGGTT	TGCAGAAAAG	350
AGCTTGATCC	ACGAGTTCAG	AAGTCCTGGT	ATGTTCTCTAG	AAAG	394
ATG TTC TCC	TGG AAA GCT	TCA AAA GCC	AGG TCT CCA	TTA AGT	436
CCA AGG TAT	TCT CTA CCT	GGT AGT ACA	GAG GTA CTT	ACA GGT	478
TGT CAT TCT	TAT CCT TCC	AGA TTC CTG	TCT GCC AGC	TCT TTT	520
ACT TCA GCC	CTG AGC ACA	GTC AAC ATG	CCT AGG GGT	CAA AAG	565
AGT AAG ACC	CGC TCC CGT	GCA AAA CGA	CAG CAG TCA	CGC AGG	604
GAG GTT CCA	GTA GTT CAG	CCC ACT GCA	GAG GAA GCA	GGG TCT	646
TCT CCT GTT	GAC CAG AGT	GCT GGG TCC	AGC TTC CCT	GGT GGT	688
TCT GCT CCT	CAG GGT GTG	AAA ACC CCT	GGA TCT TTT	GGT GCA	730
GGT GTA TCC	TGC ACA GGC	TCT GGT ATA	GGT GGT AGA	AAT GCT	772
GCT GTC CTG	CCT GAT ACA	AAA AGT TCA	GAT GGC ACC	CAG GCA	814
GGG ACT TCC	ATT CAG CAC	ACA CTG AAA	GAT CCT ATC	ATG AGG	856
AAG GCT AGT	GTG CTG ATA	GAA TTC CTG	CTA GAT AAA	TTT AAG	898
ATG AAA GAA	GCA GTT ACA	AGG AGT GAA	ATG CTG GCA	GTA GTT	940
AAC AAG AAG	TAT AAG GAG	CAA TTC CCT	GAG ATC CTC	AGG AGA	982
ACT TCT GCA	CGC CTA GAA	TTA GTC TTT	GGT CTT GAG	TTG AAG	1024
GAA ATT GAT	CCC AGC ACT	CAT TCC TAT	TTG CTG GTA	GGC AAA	1066
CTG GGT CTT	TCC ACT GAG	GGA AGT TTG	AGT AGT AAC	TGG GGG	1108
TTG CCT AGG	ACA GGT CTC	CTA ATG TCT	GTC CTA GGT	GTG ATC	1150
TTT ATG AAG	GGT AAC CGT	GCC ACT GAG	CAA GAG GTC	TGG CAA	1192
TTT CTG CAT	GGA GTG GGG	GTA TAT GCT	GGG AAG AAG	CAC TTG	1234
ATC TTT GGC	GAG CCT GAG	GAG TTT ATA	AGA GAT GTA	GTG CGG	1276
GAA AAT TAC	CTG GAG TAC	CGC CAG GTA	CCT GGC AGT	GAT CCC	1314
CCA AGC TAT	GAG TTC CTG	TGG GGA CCC	AGA GCC CAT	GCT GAA	1360
ACA ACC AAG	ATG AAA GTC	CTG GAA GTT	TTA GCT AAA	GTC AAT	1402
GGC ACA GTC	CCT AGT GCC	TTT CCT AAT	CTC TAC CAG	TTG GCT	1444
CTT AGA GAT	CAG GCA GGA	GGG GTG CCA	AGA AGG AGA	GTT CAA	1486
GGC AAG GGT	GTT CAT TCC	AAG GCC CCA	TCC CAA AAG	TCC TCT	1528
AAC ATG TAG					1537
TTGAGTCTGT	TCTGTTGTGT	TTGAAAAACA	GTCAGGCTCC	TAATCAGTAG	1587
AGAGTTTATA	GCCTACCAGA	ACCAACATGC	ATCCATTCTT	GGCCTGTTAT	1637
ACATTAGTAG	AATGGAGGCT	ATTTTGTGTA	CTTTTCAAAT	GTTTGTAA	1687
CTAAACAGTG	CTTTTTGCCA	TGCTTCTTGT	TAAGTGCATA	AAGAGGTAAC	1737
TGTCACCTGT	CAGATTAGGA	CTTGTTTTGT	TATTTGCAAC	AAACTGGAAA	1787
ACATTATTTT	GTTTTTACTA	AAACATTGTG	TAACATTGCA	TTGGAGAAGG	1837
GATTGTGATG	GCAATGTGAT	ATCATACAGT	GGTGAAACAA	CAGTGAAGTG	1887
GGAAAGTTTA	TATTGTAAAT	TTTGAAAATT	TTATGAGTGT	GATTGCTGTA	1937
TACTTTTTTC	TTTTTTGTAT	AATGCTAAGT	GAAATAAAGT	TGGATTGAT	1987
GACTTTACTC	AAATTCATTA	GAAAGTAAAT	CGTAAACTC	TATTACTTTA	2037
TTATTTCTTT	CAATTATGAA	TTAAGCATTG	GTTATCTGGA	AGTTTCTCCA	2087
GTAGCACAGG	ATCTAGTATG	AAATGTATCT	AGTATAGGCA	CTGACAGTGA	2137
GTTATCAGAG	TCT				2150

- (2) INFORMATION FOR SEQUENCE ID NO: 25:  
 (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH: 2099 base pairs  
     (B) TYPE: nucleic acid  
     (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: genomic DNA  
 (ix) FEATURE:  
     (A) NAME/KEY: smage-II  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

ACCTTATTGG	GTCTGTCTGC	ATATGCCTCC	ACTTGTGTGT	AGCAGTCTCA	50
AATGGATCTC	TCTCTACAGA	CCTCTGTCTG	TGTCTGGCAC	CCTAAGTGGC	100
TTTGCATGGG	CACAGGTTTC	TGCCCCCTGCA	TGGAGCTTAA	ATAGATCTTT	150
CTCCACAGGC	CTATACCCCT	GCATTGTAAG	TTTAAGTGGC	TTTATGTGGA	200
TACAGGTCTC	TGCCCTTGTA	TGCAGGCCTA	AGTTTTTCTG	TCTGCTTAGC	250
CCCTCCAAGT	GAAGCTAGTG	AAAGATCTAA	CCCACTTTTG	GAAGTCTGAA	300
ACTAGACTTT	TATGCAGTGG	CCTAACAAGT	TTTAATTTCT	TCCACAGGGT	350
TTGCAGAAAA	GAGCTTGATC	CACGAGTTCC	GAAGTCTCTG	TATGTTCCCTA	400
GAAAGATGTT	CTCCTGGAAA	GCTTCAAAAG	CCAGGTCTCC	ATTAAGTCCA	450
AGGTATTCTC	TACCTGGTAG	TACAGAGGTA	CTTACAGGTT	GTCATTCTTA	500
TCTTTCCAGA	TTCTGTCTG	CCAGCTCTTT	TACTTCAGCC	CTGAGCACAG	550
TCAACATGCC	TAGGGGTCAA	AAGAGTAAGA	CCCCTCCCCG	TGCAAAACGA	600
CAGCAGTCAC	GCAGGGAGGT	TCCAGTAGTT	CAGCCCACTG	CAGAGGAAGC	650
AGGGTCTTCT	CTGTTGACC	AGAGTGCTGG	GTCCAGCTTC	CCTGGTGGTT	700
CTGCTCCTCA	GGGTGTGAAA	ACCCCTGGAT	CTTTTGGTGC	AGGTGTATCC	750
TGCACAGGCT	CTGGTATAGG	TGGTAGAAAT	GCTGCTGTCC	TGCCTGATAC	800
AAAAAGTTCA	GATGGCACCC	AGGCAGGGAC	TTCCATTGAG	CACACACTGA	850
AAGATCCTAT	CATGAGGAAG	GCTAGTGTCG	TGATAGAATT	CCTGCTAGAT	900
AAGTTTAAGA	TGAAAGAAGC	AGTTACAAGG	AGTGAAATGC	TGGCAGTAGT	950
TAACAAGAAG	TATAAGGAGC	AATTCCCTGA	GATCCTCAGG	AGAACTTCTG	1000
CACGCCTAGA	ATTAGTCTTT	GGTCTTGAGT	TGAAGGAAAT	TGATCCAGC	1050
ACTCATTCCCT	ATTTGCTGGT	AGGCAAAAGT	GGTCTTTCCA	CTGAGGGAAG	1100
TTTGAGTAGT	AACTGGGGGT	TGCCTAGGAC	AGGTCTCCTA	ATGTCTGTCC	1150
TAGGTGTGAT	CTTCATGAAG	GGTAACCGTG	CCACTGAGCA	AGAGGTCTGG	1200
CAATTTCTGC	ATGGAGTGGG	GGTATATGCT	GGGAAGAAGC	ACTTGATCTT	1250
TGGCGAGCCT	GAGGAGTTTA	TAAGAGATGT	AGTGCGGGAA	AATTACCTGG	1300
AGTACCGCCA	GGTACCTGGC	AGTGATCCCC	CAAGCTATGA	GTTCTGTGG	1350
GGACCCAGAG	CCCATGCTGA	AACAACCAAG	ATGAAAGTCC	TGGAAGTTTT	1400
AGCTAAAGTC	AATGGCACAG	TCCCTAGTGC	CTTCCCTAAT	CTCTACCAGT	1450
TGGCTCTTAG	AGATCAGGCA	GGAGGGGTGC	CAAGAAGGAG	AGTTCAAGGC	1500
AAGGGTGTTC	ATTCCAAGGC	CCCATCCCAA	AAGTCCCTCTA	ACATGTAGTT	1550
GAGTCTGTTC	TGTTGTGTTT	GAAAAACAGT	CAGGCTCCTA	ATCAGTAGAG	1600
AGTTTCATAGC	CTACCAGAAC	CAACATGCAT	CCATTCTTGG	CCTGTTATAC	1650
ATTAGTAGAA	TGGAGGCTAT	TTTGTGTTACT	TTTCAAATGT	TTGTTTAACT	1700
AAACAGTGCT	TTTGGCCATG	CTTCTTGTTA	ACTGCATAAA	GAGGTAAGT	1750
TCACCTTGTC	GATTAGGACT	TGTTTGTGTTA	TTTGCAACAA	ACTGGAAAAC	1800
ATTATTTTGT	TTTTACTAAA	ACATTGTGTA	ACATTGCATT	GGAGAAGGGA	1850
TTGTCATGGC	AATGTGATAT	CATACAGTGG	TGAAACAACA	GTGAAGTGGG	1900
AAAGTTTATA	TTGTTAGTTT	TGAAAATTTT	ATGAGTGTGA	TTGCTGTATA	1950
CTTTTTTCTT	TTTGTATAA	TGCTAAGTGA	AATAAAGTTG	GATTTGATGA	2000
CTTTACTCAA	ATTCATTAGA	AAGTAAATCA	TAAAACTCTA	TTACTTTATT	2050
ATTTTCTTCA	ATTATTAATT	AAGCATTGGT	TATCTGGAAG	TTTCTCCAG	2099

- (2) INFORMATION FOR SEQUENCE ID NO: 26:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 9 amino acids
    - (B) TYPE: amino acids
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: protein
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Glu Ala Asp Pro Thr Gly His Ser Tyr  
5

We claim:

1. Isolated nucleic acid molecule which codes for or is complementary to a nucleic acid molecule which codes for tumor rejection antigen precursor MAGE-3.
2. The isolated nucleic acid molecule of claim 1, which codes for tumor rejection antigen precursor MAGE-3.
3. The nucleic acid molecule of claim 2, wherein said molecule is cDNA.
4. The nucleic acid molecule of claim 3, wherein said molecule has the nucleotide sequence set forth in SEQ ID NO: 1 (MAGE-3) or SEQ ID NO: 2 (MAGE-31).
5. Expression vector comprising the nucleic acid molecule of claim 2 operably linked to a promoter.
6. The expression vector of claim 5, further comprising a nucleic acid molecule which codes for HLA-A1.
7. Cell line transfected with the nucleic acid molecule of claim 2.
8. The cell line of claim 7, wherein said cell line expresses HLA-A1.
9. The cell line of claim 7, wherein said cell line is further transfected with a nucleic acid molecule which codes for HLA-A1.
10. Isolated tumor rejection antigen precursor coded for by the nucleic acid molecule of claim 2.
11. Vaccine comprising the isolated tumor rejection antigen precursor of claim 10 and an adjuvant.

12. Isolated tumor rejection antigen derived from the tumor rejection antigen precursor of claim 10, wherein said tumor rejection antigen is antigen D.

5 13. Isolated complex of the tumor rejection antigen of claim 12 and HLA-A1.

10 14. Method for treating a disorder characterized by expression of tumor rejection antigen precursor MAGE-3, comprising administering to a subject an amount of a cytolytic T cell specific for complexes of a tumor rejection antigen derived from MAGE-3 and a human leukocyte antigen molecule, sufficient to generate an immune response against said complexes.

15 15. The method of claim 14, wherein said human leukocyte antigen is HLA-A1.

20 16. The method of claim 15, wherein said tumor rejection antigen is antigen D.

25 17. Method for treating a disorder characterized by expression of tumor rejection antigen precursor MAGE-3, comprising administering an agent sufficient to provoke an immune response to complexes of a tumor rejection antigen derived from MAGE-3 and a human leukocyte antigen, to a subject in need thereof.

30 18. The method of claim 17, wherein said human leukocyte antigen is HLA-A1.


35 19. The method of claim 18, wherein said tumor rejection antigen is antigen D.

20. Method for determining a disorder characterized by expression of tumor rejection antigen precursor MAGE-3, comprising contacting a sample taken from a subject with an agent which identifies said tumor rejection antigen

precursor to determine expression of said tumor rejection antigen precursor as a determination of said disorder.


- 5           21. Method for determining a disorder characterized by  
expression of tumor rejection antigen precursor MAGE-3  
and presentation of a tumor rejection antigen derived  
therefrom by a cell, comprising contacting a sample taken  
with a subject with an agent which identifies said tumor  
10 rejection antigen to determine said tumor rejection  
antigen as a determination of said disorder.

1/18

**FIG. 1A**

2/18


FIG. 1B






3/18


FIG. 2



4/18


**FIG. 3**

5 / 18  
**FIG. 4**




6/18



FIG. 5



7/18

**FIG. 6**

8/18

**FIG. 7****FIG. 8**


## FIG. 9

MAGE-3 /// CCTCCCCAGAGTCTCTCAGGGAGCCTTCagCctTcCCCCACTACCATgAACtaCccCTCcttgGAGccCAAtCCtTaTGAGGacTCCAGCAaCCaaGAAGAGGAGG  
 MAGE-2 // CCTCCCCAcAGTCTCTCAGGGAGCCTTCagCTTtctCGACTACCATCAACTaCACtCTctttgGAGaCAAtCCCGaTGAGGGcTCCAGCAaCCaaGAAGAGGAGG  
 MAGE-1 / CCTCCCCAGAGTCTCTCAGGGAGCCTTCGGCCtTTCCCACTACCAATCAACTtCACTtCGACAGAGGGCAACCCAGtGAGGGtTCCAGCAAGCCCTGtGAAGAGGAGG  
 225 CHO-8  
 /// GGCCAAAGCAGCCTtcccTgaCC - TGGAGTCCgaGTTCCCaAGCAGcAcTCAgTAGAAGGTGGCCcGAGtTGGTtTcaATTtTCTGCTCTCTCAAgTATTCGAGGCCA  
 // GGCCAAAGaAtgTtTcccgaCCtTGGAGTCCCGAGTTCaAGCAGcAAATCAgTAGGAAGaTGgTtGAGtTGGTtTcaATTtTCTGCTCTCTCAAgTATTCGAGGCCA  
 / GGCCAAAGCAGCCTtCTtGtATCC - TGGAGTCCtTTGtTTCGGAGCAGTAATCACTAAgAGGTGGCTGAATTtTGGTtGgTtTCTGCTCTCAAAATATTCGAGGCCA  
 325  
 /// GGAGGCCgGTTCACAAAGGCAGAAATGCTGGgGAGTGTCTgTcgAAATtTggcAGtAtTtctTTCTCTGtGATCTTcAGCAAAAGcLTCCagTTCCTTGCAGCT  
 // GGAGGCCgGTTCACAAAGGCAGAAATGCTGGAGAGTGTCTcTCAGAAATtTgcAGgACTtctTTTCCcGtGATCTTcAGCAAAAGCCCTCCGAGTACTTGCAGCT  
 / GGAGGCCAGTCAcAAAGGCAGAAATGCTGGAGAGTGTCAITCAAAATtTAcAAGcACTGtTtTCTCTGAGATCTTTCGGCAAAAGCCCTtGAGtTCTTGCAGCT  
 425 SEQ-4  
 /// GGTCtTTTGGCATcGAgcTGATGGAAgtGGACCCcATcGGGCCACTtGtTAcATCTtTTGccACCCTGCCCTgGGccCTCTCTCTA cGATGGCCCTGCTGGGTGAcAAT  
 // GGTCtTTTGGCATcGAgGTGgtGGAAgtGGtCCCCATcAgGCCACTtGtTAcATCTCTTGTACCTTGCCCTgGGcCTCTCTCTAGATGGCCCTGCTGGGCGAcAAT  
 / GGTCtTTTGGCATtGACGTGAAGGAAGCAGACCCcACCGCCACtTCCtATtGTCCTtGTCACCTtGCCCTAGGTCTCTCTTAATGATGGCCCTGCTGGGTGATAT  
 525  
 /// CAGATCATGCCCCAAGgCAGGCCcTCTCTGATAATcGTCTTGGcCATaATcGCAAGaGAGGGCGGAcTgTGCcCTTGAGGAGaAAATCTTGGGAGGAGCTGAGTG  
 // CAGtTCATGCCCCAAGACAGGCCcTCTCTGATAATcCTC - TGGcCATaATcGCAATaGAGGGCGGAcTgTGCcCTCTGAGGAGaAAATCTTGGGAGGAGCTGAGTa  
 / CAGATCATGCCCCAAGcAGGCtTCTCTGATaATtGTCCTCTGTCATGATtGCAATGGAGGGCGGCCATCTCTCTGAGGAGGAATCTTGGGAGGAGCTGAGTG  
 625 CHO-9


10/18

 $\beta$ -action

MAGE


PROBES

MZ2-MEL.3.0  
MZ2-MEL 1982  
MZ2-MEL.2.2 E-  
MZ2-PBL-PHA  
Lung  
Kidney



MZ2-MEL 3.0  
MZ2-CTL 82/30  
LB34-MEL  
LB17-MEL  
MI665/2-MEL  
LB41-MEL  
MI10221-MEL  
MI13443-MEL  
SK23-MEL  
SK33-MEL

Other  
melanomas



LB4-MEL  
MI4024-MEL  
MZ3-MEL  
MZ5-MEL  
SK29-MEL  
LB31-COL  
LS411-COL



H209-SCLC  
H345-SCLC  
H510-SCLC  
TT

Other  
tumors

FIG. 10



11/18

FIG. 11A

		EXPRESSION OF MAGE GENE FAMILY				RECOGNITION BY ANI-E CTL		Expression of antigen MZ2-E after transfection**	
		Northern blot probed with cross-reactive MAGE-1 probe*	cDNA-PCR product probed with oligonucleotide specific for:			tested by:			
			MAGE-1	MAGE-2	MAGE-3†	TNF release‡	Lysis§		
Cells of patient MZ2	melanoma cell line MZ2-MEL.3.0	+	+++	+++	+++	+	+		
	tumor sample MZ2 (1982)	+	+++	+++	+++				
	antigen-loss variant MZ2-MEL.2.2	+	-	+++	+++	-	-		
	CTL clone MZ2-CTL.82/30	-	-	-	-				
	PHA-activated blood lymphocytes	-	-	-	-				
Normal tissues	Liver	-	-	-	-				
	Muscle	-	-	-	-				
	Skin	-	-	-	-				
	Lung	-	-	-	-				
	Brain	-	-	-	-				
	Kidney	-	-	-	-				
Melanoma cell lines of HLA-A1 patients	LB34-MEL	+	++	+++	+++	+	+	+	
	MI665/2-MEL	-	-	-	-	-	-	+	
	MI10221-MEL	+	-	++	+++	-	-		
	MI13443-MEL	+	++	+++	+++	+	+		
	SK33-MEL	+	-	+++	+++	-	-	-	
	SK23-MEL	+	-	+++	+++	-	-	+	

\* Data obtained in the conditions of figure 5.

† Data obtained as described in figure 6.

‡ TNF release by CTL 82/30 after stimulation with the tumor cells as described in (11).

§ Lysis of 51 Cr labeled target by CTL 82/30 in the conditions of figure 1.

\*\* Cells transfected with the 2.4 kb fragment of gene MAGE-1 were tested for their ability to stimulate TNF release by CTL 82/30

FIG. 11B

12/18							
Melanoma cell lines of other patients	LB17-MEL	+	+	+	++	++	-
	LB33-MEL	+	-	+	++	++	-
	LB4-MEL	-	-	-	-	-	-
	LB41-MEL	-	-	-	-	-	-
	MI4024-MEL	+	++	+	++	++	-
	SK29-MEL	-	-	-	-	-	-
	MZ3-MEL	+	+	+	++	++	-
	MZ5-MEL	+	-	+	++	++	-
	Melanoma tumor sample	BB5-MEL	+	++	+	++	++
Other tumor cell lines	small cell lung cancer H209	+	-	+	++	++	-
	small cell lung cancer H345	+	-	+	++	++	-
	small cell lung cancer H510	+	-	+	++	++	-
	small cell lung cancer LB11	+	+	+	++	++	-
	bronchial squamous cell carcinoma LB37	+	-	+	++	++	-
	thyroid medullary carcinoma TT	+	++	+	++	++	-
	colon carcinoma LB31	+	-	+	++	++	-
	colon carcinoma LS411	-	-	-	-	-	-
	Other tumor samples	chronic myeloid leukemia LLC5	-	-	-	-	-
acute myeloid leukemia TA		-	-	-	-	-	-

12/18

\* Data obtained in the conditions of figure 5.

† Data obtained as described in figure 6.

‡ TNF release by CTL 82/30 after stimulation with the tumor cells as described in (11).

§ Lysis of 51 Cr labelled target by CTL 82/30 in the conditions of figure 1.

\*\* Cells transfected with the 2.4 kb fragment of gene MAGE-1 were tested for their ability to stimulate TNF release by CTL 82/30

13 / 18

**FIG. 12**





FIG. 13



15/18

FIG. 14B









FIG. 14A



16/18

**FIG. 15A****FIG. 15B**

17/18

**FIG. 15C****FIG. 15D**

18/18




FIG. 16



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/02877

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) : Please See Extra Sheet.

US CL : 935/12, 55, 70; 530/350, 395, 828; 424/88; 435/6, 7.1

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 935/12, 55, 70; 530/350, 395, 828; 424/88; 435/6, 7.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, BIOSIS, MEDLINE, EMBASE, CA

search terms: antigen D, MAGE?

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	SCIENCE, Volume 254, issued 13 December 1991, Van der Bruggen et al., "A Gene Encoding an Antigen Recognized by Cytolytic T Lymphocytes on a Human Melanoma", pages 1643-1647, see entire document.	1-4 ---- 5-21
Y	INTERNATIONAL JOURNAL OF CANCER, Volume 52, issued 1992, Brassuer et al., "Human Gene MAGE-1, Which Codes For a Tumor Rejection Antigen, Is Expressed by Some Breast Tumors", pages 839-841, see entire document.	11-21

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be part of particular relevance	"T"	later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention
* "E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* "L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* "O"	document referring to an oral disclosure, use, exhibition or other means	* "A"	document member of the same patent family
* "P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

14 MAY 1994

Date of mailing of the international search report

JUN 16 1994

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ANTHONY C. CAPUTA

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/02877

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	IMMUNOGENETICS, Volume 35, issued 1992, Traversari et al., "Transfection and Expression of a Gene Coding for a Human Melanoma Antigen Recognized by Autologous Cytolytic T Lymphocytes", pages 145-152, see entire document.	5-21
Y	INTERNATIONAL JOURNAL OF CANCER, Volume 39, issued 1987, M. Herin et al., "Production of Stable Cytolytic T-Cell Clones Directed Against Autologous Human Melanoma", pages 390-396, see entire document.	5-21

# INTERNATIONAL SEARCH REPORT

Int. application No.

PCT/US94/02877

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (5):

C12N 15/12, 15/85; C07K 15/06, 15/14; C12Q 1/68; G01N 33/53; A61K 39/00,

